




Land Cover

Summarized by: **Big Horn** (County)



31%
(1,006,890
Acres)

Grassland Systems Lowland/Prairie Grassland

Great Plains Mixedgrass Prairie

The system covers much of the eastern two-thirds of Montana, occurring continuously for hundreds of square kilometers, interrupted only by wetland/riparian areas or sand prairies. Soils are primarily fine and medium-textured. The growing season averages 115 days, ranging from 100 days on the Canadian border to 130 days on the Wyoming border. Climate is typical of mid-continental regions with long severe winters and hot summers. Grasses typically comprise the greatest canopy cover, and western wheatgrass (*Pascopyrum smithii*) is usually dominant. Other species include thickspike wheatgrass (*Elymus lanceolatus*), green needlegrass (*Nassella viridula*), blue grama (*Bouteloua gracilis*), and needle and thread (*Hesperostipa comata*). Near the Canadian border in north-central Montana, this system grades into rough fescue (*Festuca campestris*) and Idaho fescue (*Festuca idahoensis*) grasslands. Remnants of shortbristle needle and thread (*Hesperostipa curtisetata*) dominated vegetation are found in northernmost Montana and North Dakota, and are associated with productive sites, now mostly converted to farmland. Forb diversity is typically high. In areas of southeastern and central Montana where sagebrush steppe borders the mixed grass prairie, common plant associations include Wyoming big sagebrush-western wheatgrass (*Artemisia tridentata* ssp. *wyomingensis*/ *Pascopyrum smithii*). Fire and grazing are the primary drivers of this system. Drought can also impact it, in general favoring the shortgrass component at the expense of the mid-height grasses. With intensive grazing, cool season exotics such as Kentucky bluegrass (*Poa pratensis*), smooth brome (*Bromus inermis*), and Japanese brome (*Bromus japonicus*) increase in dominance; both of these rhizomatous species have been shown to markedly decrease species diversity. Previously cultivated acres that have been re-vegetated with non-native plants have been transformed into associations such as Kentucky bluegrass (*Poa pratensis*)/western wheatgrass (*Pascopyrum smithii*) or into pure crested wheatgrass (*Agropyron cristatum*) stands.



18%
(568,391
Acres)

Shrubland, Steppe and Savanna Systems

Sagebrush Steppe

Big Sagebrush Steppe

This widespread ecological system occurs throughout much of central Montana, and north and east onto the western fringe of the Great Plains. In central Montana, where this system occurs on both glaciated and non-glaciated landscapes, it differs slightly, with more summer rain than winter precipitation and more precipitation annually. Throughout its distribution, soils are typically deep and non-saline, often with a microphytic crust. This shrub-steppe is dominated by perennial grasses and forbs with greater than 25% cover. Overall shrub cover is less than 10 percent. In Montana and Wyoming, stands are more mesic, with more biomass of grass, and have less shrub diversity than stands farther to the west, and 50 to 90% of the occurrences are dominated by Wyoming big sagebrush with western wheatgrass (*Pascopyrum smithii*). Japanese brome (*Bromus japonicus*) and cheatgrass (*Bromus tectorum*) are indicators of disturbance, but cheatgrass is typically not as abundant as in the Intermountain West, possibly due to a colder climate. The natural fire regime of this ecological system maintains a patchy distribution of shrubs, preserving the steppe character. Shrubs may increase following heavy grazing and/or with fire suppression. In central and eastern Montana, complexes of prairie dog towns are common in this ecological system.



11%
(339,458
Acres)

Human Land Use

Agriculture

Cultivated Crops

These areas used for the production of crops, such as corn, soybeans, small grains, sunflowers, vegetables, and cotton, typically on an annual cycle. Agricultural plant cover is variable depending on season and type of farming. Other areas include more stable land cover of orchards and vineyards.



9%
(339,458
Acres)

Forest and Woodland Systems

Conifer-dominated forest and woodland (xeric-mesic)

Great Plains Ponderosa Pine Woodland and Savanna

These ponderosa pine (*Pinus ponderosa*) occurrences differ from the Rocky Mountain Ponderosa Pine Woodland and Savanna systems in that they are typically found within the matrix of the Great Plains grassland systems. They are often surrounded by mixed-grass prairie, in places where available soil moisture is higher or soils are more coarse and rocky. Elevation ranges from 1,189 meters (3,900 feet) in southeastern Montana to 1,646 m (5,400 feet) in north-central Montana. Occurrences are usually on east- and north-facing aspects. These woodlands can be physiognomically variable, ranging from very sparse patches of trees on drier sites, to nearly closed-canopy forest stands on north slopes or in draws where available soil moisture is higher.



4%
(137,387
Acres)

Grassland Systems

Montane Grassland

Rocky Mountain Lower Montane, Foothill, and Valley Grassland

This grassland system of the northern Rocky Mountains is found at lower montane to foothill elevations in mountains and valleys throughout Montana. These grasslands are floristically similar to Big Sagebrush Steppe but are defined by shorter summers, colder winters, and young soils derived from recent glacial and alluvial material. They are found at elevations from 548 - 1,650 meters (1,800-5,413 feet). In the lower montane zone, they range from small meadows to large open parks surrounded by conifers; below the lower treeline, they occur as extensive foothill and valley grasslands. Soils are relatively deep, fine-textured, often with coarse fragments, and non-saline. Microphytic crust may be present in high-quality occurrences. This system is typified by cool-season perennial bunch grasses and forbs (>25%) cover, with a sparse shrub cover (<10%). Rough fescue (*Festuca campestris*) is dominant in the northwestern portion of the state and Idaho fescue (*Festuca idahoensis*) is dominant or co-dominant throughout the range of the system. Bluebunch wheatgrass (*Pseudoroegneria spicata*) occurs as a co-dominant throughout the range as well, especially on xeric sites. Western wheatgrass (*Pascopyrum smithii*) is consistently present, often with appreciable coverage (>10%) in lower elevation occurrences in western Montana and virtually always present, with relatively high coverages (>25%), on the edge of the Northwestern Great Plains region. Species diversity ranges from a high of more than 50 per 400 square meter plot on mesic sites to 15 (or fewer) on xeric and disturbed sites. Most occurrences have at least 25 vascular species present. Farmland conversion, noxious species invasion, fire suppression, heavy grazing and oil and gas development are major threats to this system.



Wetland and Riparian Systems Floodplain and Riparian

4%
(136,648
Acres)

Great Plains Riparian

This system is associated with perennial to intermittent or ephemeral streams throughout the northwestern Great Plains. In Montana, it occurs along smaller tributaries of the Yellowstone and Missouri rivers, as well as tributaries to the large floodplain rivers that feed them (e.g. the Milk, Marias, Musselshell, Powder, Clark's Fork Yellowstone, Tongue, etc). In areas adjacent to the mountain ranges of central and southeastern Montana, and near the Rocky Mountain Front, it grades into Rocky Mountain Lower Montane-Foothill Riparian Woodland and Shrubland systems. This system is found on alluvial soils in highly variable landscape settings, from confined, deep cut ravines to wide, braided streambeds. Channel migration occurs in less-confined areas, but within a more narrow range than would occur in broad, alluvial floodplains. Typically, the rivers are wadeable by mid-summer.

The primary inputs of water to these systems include groundwater discharge, overland flow, and subsurface interflow from the adjacent upland. Flooding is the key ecosystem process, creating suitable sites for seed dispersal and seedling establishment, and controlling vegetation succession. Communities within this system range from riparian forests and shrublands to tallgrass wet meadows and gravel/sand flats. Dominant species are similar to those found in the Great Plains Floodplain System. In the western part of the system's range in Montana, the dominant overstory species is black cottonwood (*Populus balsamifera ssp. trichocarpa*) with narrowleaf cottonwood (*Populus angustifolia*) and Plains cottonwood (*Populus deltoides*) occurring as co-dominants in the riparian/floodplain interface near the mountains. Further east, narrowleaf cottonwood and Plains cottonwood become dominant. In wetter systems, the understory is typically willow (*Salix spp.*) and redosier dogwood (*Cornus stolonifera*) with graminoids such as western wheatgrass (*Pascopyrum smithii*) and forbs like American licorice (*Glycyrrhiza lepidota*). In areas where the channel is incised, the understory may be dominated by big sagebrush (*Artemisia tridentata*) or silver sagebrush (*Artemisia cana*). Like floodplain systems, riparian systems are often subjected to overgrazing and/or agriculture and can be heavily degraded, with salt cedar (*Tamarix ramosissima*) and Russian olive (*Eleagnus angustifolia*) replacing native woody vegetation and regrowth. Groundwater depletion and lack of fire have resulted in additional species changes.



4%
(131,390
Acres)

Shrubland, Steppe and Savanna Systems Sagebrush Steppe

Montane Sagebrush Steppe

This system dominates the montane and subalpine landscape of southwestern Montana from valley bottoms to subalpine ridges and is found as far north as Glacier National Park. It can also be seen in the island mountain ranges of the north-central and south-central portions of the state. It primarily occurs on deep-soiled to stony flats, ridges, nearly flat ridgetops, and mountain slopes. In general, this system occurs in areas of gentle topography, fine soils, subsurface moisture or mesic conditions, within zones of higher precipitation and areas of snow accumulation. It occurs on all slopes and aspects, variable substrates and all soil types. The shrub component of this system is generally dominated by mountain big sagebrush (*Artemisia tridentata ssp. vaseyana*). Other co-dominant shrubs include silver sagebrush (*Artemisia cana ssp. viscidula*), subalpine big sagebrush (*Artemisia tridentata ssp. spiciformis*), three tip sagebrush (*Artemisia tripartita ssp. tripartita*) and antelope bitterbrush (*Purshia tridentata*). Little sagebrush (*Artemisia arbuscula ssp. arbuscula*) shrublands are only found in southwestern Montana on sites with a perched water table. Wyoming big sagebrush (*Artemisia tridentata ssp. wyomingensis*) sites may be included within this system if occurrences are at montane elevations, and are associated with montane graminoids such as Idaho fescue (*Festuca idahoensis*), spike fescue (*Leucopoa kingii*), or poverty oatgrass (*Danthonia intermedia*). In areas where sage has been eliminated by human activities like burning, disking or poisoning, other shrubs may be dominant, especially rubber rabbitbrush (*Ericameria nauseosa*), and green rabbitbrush (*Chrysothamnus viscidiflorus*). Because of the mesic site conditions, most occurrences support a diverse herbaceous undergrowth of grasses and forbs. Shrub canopy cover is extremely variable, ranging from 10 percent to as high as 40 or 50 percent.



2% (58,226
Acres)

Sparse and Barren Systems Bluff, Badland and Dune

Great Plains Badlands

The Western Great Plains Badlands ecological system occurs within the mixed grass and sand prairie regions of eastern and southeastern Montana, where the land lies well above or below its local base level, shaped by the carving action of streams, erosion, and erodible parent material. It is easily recognized by its rugged, eroded, and often colorful land formations, and the relative absence of vegetative cover. In those areas with vegetation, species can include scattered individuals of many dryland shrubs or herbaceous taxa, including curlycup gumweed (*Grindelia squarrosa*), threadleaf snakeweed (*Gutierrezia sarothrae*) (especially with overuse and grazing), greasewood (*Sarcobatus vermiculatus*), Gardner's saltbush (*Atriplex gardneri*), buckwheat (*Eriogonum* species), plains muhly (*Muhlenbergia cuspidata*), bluebunch wheatgrass (*Pseudoroegneria spicata*), and Hooker's sandwort (*Arenaria hookeri*). Patches of sagebrush (*Artemisia* spp.) can also occur. Climate is typical of mid continental regions with long severe winters and warm summers. Precipitation ranges from 7 to 14 inches per year, with two-thirds of the precipitation falling during the summer, and a third falling in the spring. The sedimentary parent material of exposed rocks and the resultant eroded clay soils are derived from Cretaceous sea beds and are often fossil-rich. Dominant soil types are in the order Entisols. These mineral soils are found primarily on uplands, slopes, and creek bottoms and are easily erodible. The growing season is short, averaging 115 days, with a range from 100 days on the Canadian border to 130 days on the Wyoming border. Land use is limited, except for off-highway vehicle recreation and incidental grazing.



2% (54,176
Acres)

Human Land Use Agriculture

Pasture/Hay

These agriculture lands typically have perennial herbaceous cover (e.g. regularly-shaped plantings) used for livestock grazing or the production of hay. There are obvious signs of management such as irrigation and haying that distinguish it from natural grasslands. Identified CRP lands are included in this land cover type.



Grassland Systems Lowland/Prairie Grassland

Great Plains Sand Prairie

2% (53,492 Acres)

The sand prairies constitute a very unique system within the western Great Plains. The unifying and controlling feature for this system is that coarse-textured soils predominate and the dominant grasses are well-adapted to this condition. In the northwestern portion of the system's range, stand size corresponds to the area of exposed caprock sandstone, and small patches predominate, but larger patches are found embedded in the encompassing Great Plains Mixed Grass Prairie, and usually occupy higher positions in local landscapes where former caprock formations have eroded into more subdued and planar topography. In most of eastern Montana, substrates supporting this system have weathered in place from sandstone caprock. Soils can be relatively thin or deep due to varying amounts of downslope movement of weathered sands. Needle and thread (*Hesperostipa comata*) is the dominant grass species. Other frequent species include little bluestem (*Schizachyrium scoparium*), often occurring with threadleaf sedge (*Carex filifolia*) and dominating both sandy sites and actively eroding sites. Prairie sandreed (*Calamovilfa longifolia*), sand bluestem (*Andropogon hallii*) and big bluestem (*Andropogon gerardii*) are sporadically distributed and found generally on the coarsest-textured sands. Other graminoids include bluebunch wheatgrass (*Pseudoroegneria spicata*), sun sedge (*Carex inops ssp. heliophila*), and purple threeawn (*Aristida purpurea*). Characteristic forbs differ by occurrence, but species of scurf pea (*Psoralegium* species) and Indian breadroot (*Pediemelum*) species are common. Communities of silver sage (*Artemisia cana ssp. cana*) or skunkbush sumac (*Rhus trilobata*) can occur within this system. Wind erosion, fire and grazing constitute the other major dynamic processes that can influence this system.



Forest and Woodland Systems Conifer-dominated forest and woodland (xeric-mesic)

Rocky Mountain Dry-Mesic Montane Mixed Conifer Forest

2% (52,970 Acres)

This ecological system, composed of highly variable montane conifer forests, is found throughout Montana. It is associated with a submesic climate regime with annual precipitation ranging from 250 to 1,000 millimeters (10-39 inches), with most precipitation occurring during winter, and April through June. Winter snowpacks typically melt off in early spring at lower elevations. Elevations range from valley bottoms to 1,676 meters (5,500 feet) in northwestern Montana and up to 2,286 meters (7,500 feet) on warm aspects in southern Montana. In northwestern and west-central Montana, this ecosystem forms a forest belt on warm, dry to slightly moist sites. It generally occurs on gravelly soils with good aeration and drainage and a neutral to slightly acidic pH. In the western part of the state, it is seen mostly on well drained mountain slopes and valleys from lower treeline to up to 1,676 meters (5,500 feet). Immediately east of the Continental Divide, in north-central Montana, it occurs at montane elevations. Douglas-fir (*Pseudotsuga menziesii*) is the dominant conifer both as a seral and climax species. West of the Continental Divide, occurrences can be dominated by any combination of Douglas-fir and long-lived, seral western larch (*Larix occidentalis*), grand fir (*Abies grandis*), ponderosa pine (*Pinus ponderosa*) and lodgepole pine (*Pinus contorta*). Aspen (*Populus tremuloides*) and western white pine (*Pinus monticola*) have a minor status, with western white pine only in extreme western Montana. East of the Continental Divide, larch is absent and lodgepole pine is the co-dominant. Engelmann spruce (*Picea engelmannii*), white spruce, (*Picea glauca*) or their hybrid, become increasingly common towards the eastern edge of the Douglas-fir forest belt.

Additional Limited Land Cover

- 1% (42,114 Acres) ■ [Recently burned forest](#)
- 1% (36,768 Acres) ■ [Recently burned shrubland](#)
- 1% (36,430 Acres) ■ [Burned Sagebrush](#)
- 1% (34,610 Acres) ■ [Rocky Mountain Montane Douglas-fir Forest and Woodland](#)
- 1% (28,380 Acres) ■ [Recently burned grassland](#)
- 1% (22,515 Acres) ■ [Post-Fire Recovery](#)
- 1% (22,096 Acres) ■ [Rocky Mountain Foothill Limber Pine - Juniper Woodland](#)
- 1% (20,096 Acres) ■ [Other Roads](#)
- 1% (19,360 Acres) ■ [Great Plains Floodplain](#)
- 1% (17,605 Acres) ■ [Great Plains Wooded Draw and Ravine](#)
- <1% (15,534 Acres) ■ [Greasewood Flat](#)
- <1% (13,925 Acres) ■ [Quarries, Strip Mines and Gravel Pits](#)
- <1% (10,659 Acres) ■ [Open Water](#)
- <1% (7,712 Acres) ■ [Rocky Mountain Lodgepole Pine Forest](#)
- <1% (7,457 Acres) ■ [Mountain Mahogany Woodland and Shrubland](#)
- <1% (6,616 Acres) ■ [Great Plains Cliff and Outcrop](#)
- <1% (6,427 Acres) ■ [Introduced Upland Vegetation - Annual and Biennial Forbland](#)
- <1% (5,132 Acres) ■ [Introduced Upland Vegetation - Perennial Grassland and Forbland](#)
- <1% (5,088 Acres) ■ [Introduced Riparian and Wetland Vegetation](#)
- <1% (4,602 Acres) ■ [Interstate](#)
- <1% (4,391 Acres) ■ [Aspen Forest and Woodland](#)
- <1% (3,380 Acres) ■ [Low Intensity Residential](#)
- <1% (3,269 Acres) ■ [Major Roads](#)

<1% (3,152 Acres) ■ [Rocky Mountain Subalpine Dry-Mesic Spruce-Fir Forest and Woodland](#)

<1% (2,928 Acres) ■ [Mat Saltbush Shrubland](#)

<1% (1,785 Acres) ■ [Railroad](#)

<1% (1,494 Acres) ■ [Rocky Mountain Lower Montane-Foothill Shrubland](#)

<1% (1,490 Acres) ■ [Great Plains Saline Depression Wetland](#)

<1% (989 Acres) ■ [Developed, Open Space](#)

<1% (883 Acres) ■ [Great Plains Closed Depressional Wetland](#)

<1% (681 Acres) ■ [Low Sagebrush Shrubland](#)

<1% (674 Acres) ■ [Commercial / Industrial](#)

<1% (576 Acres) ■ [Rocky Mountain Subalpine-Upper Montane Grassland](#)

<1% (549 Acres) ■ [Introduced Upland Vegetation - Annual Grassland](#)

<1% (502 Acres) ■ [Great Plains Open Freshwater Depression Wetland](#)

<1% (398 Acres) ■ [Rocky Mountain Cliff, Canyon and Massive Bedrock](#)

<1% (341 Acres) ■ [Emergent Marsh](#)

<1% (331 Acres) ■ [Big Sagebrush Shrubland](#)

<1% (246 Acres) ■ [Rocky Mountain Montane-Foothill Deciduous Shrubland](#)

<1% (230 Acres) ■ [High Intensity Residential](#)

<1% (113 Acres) ■ [Rocky Mountain Subalpine-Montane Fen](#)

<1% (70 Acres) ■ [Rocky Mountain Subalpine-Montane Riparian Shrubland](#)

<1% (61 Acres) ■ [Coal Bed Methane](#)

<1% (57 Acres) ■ [Rocky Mountain Subalpine-Montane Mesic Meadow](#)

<1% (37 Acres) ■ [Aspen and Mixed Conifer Forest](#)

<1% (26 Acres) ■ [Gas and Gas Storage](#)

<1% (13 Acres) ■ [Oil and Oil / Gas](#)

<1% (7 Acres) ■ [Great Plains Shrubland](#)

<1% (6 Acres) ■ [Introduced Upland Vegetation - Shrub](#)

<1% (3 Acres) ■ [Injection](#)

<1% (2 Acres) ■ [Mixed Salt Desert Scrub](#)

<1% (0 Acres) ■ [Northern Rocky Mountain Lower Montane Riparian Woodland and Shrubland](#)

Introduction to Land Cover

Land Use/Land Cover is one of 15 [Montana Spatial Data Infrastructure](#) framework layers considered vital for making statewide maps of Montana and understanding its geography. The layer records all Montana natural vegetation, land cover and land use, classified from satellite and aerial imagery, mapped at a scale of 1:100000, and interpreted with supporting ground-level data. The baseline map is adapted from the Northwest ReGAP (NWGAP) project land cover classification, which used 30m resolution multi-spectral Landsat imagery acquired between 1999 and 2001. Vegetation classes were drawn from the Ecological System Classification developed by NatureServe (Comer et al. 2003). The land cover classes were developed by Anderson et al. (1976). The NWGAP effort encompasses 12 map zones. Montana overlaps seven of these zones. The two NWGAP teams responsible for the initial land cover mapping effort in Montana were Sanborn and NWGAP at the University of Idaho. Both Sanborn and NWGAP employed a similar modeling approach in which Classification and Regression Tree (CART) models were applied to Landsat ETM+ scenes. The Spatial Analysis Lab within the Montana Natural Heritage Program was responsible for developing a seamless Montana land cover map with a consistent statewide legend from these two separate products. Additionally, the Montana land cover layer incorporates several other land cover and land use products (e.g., MSDI Structures and Transportation themes and the Montana Department of Revenue Final Land Unit classification) and reclassifications based on plot-level data and the latest NAIP imagery to improve accuracy and enhance the usability of the theme. Updates are done as partner support and funding allow, or when other MSDI datasets can be incorporated. Recent updates include fire perimeters and agricultural land use (annually), energy developments such as wind, oil and gas installations (2014), roads, structures and other impervious surfaces (various years): and local updates/improvements to specific ecological systems (e.g., central Montana grassland and sagebrush ecosystems). Current and previous versions of the Land Use/Land Cover layer with full metadata are available for download at the Montana State Library's [Geographic Information Clearinghouse](#).

Within the report area you have requested, land cover is summarized by acres of Level 1, Level 2, and Level 3 Ecological Systems.

Literature Cited

- Anderson, J.R. E.E. Hardy, J.T. Roach, and R.E. Witmer. 1976. A land use and land cover classification system for use with remote sensor data. U.S. Geological Survey Professional Paper 964.
- Comer, P., D. Faber-Langendoen, R. Evans, S. Gawler, C. Josse, G. Kittel, S. Menard, M. Pyne, M. Reid, K. Schulz, K. Snow, and J. Teague. 2003. Ecological systems of the United States: A working classification of U.S. terrestrial systems. NatureServe, Arlington, VA.