

County	Yellowstone	Upstream River Mile	362.2
Classification	UB: Unconfined braided	Downstream River Mile	357.9
General Location	East Billings	Length	4.30 mi (6.92 km)
General Comments	Wide corridor d/s Billings; WAI Reach F		

Narrative Summary

Reach B3 is 4.3 miles long and located in east Billings. The reach is characterized by the loss of several miles of side channel, extensive Russian olive infestation, and substantial flow alterations due to human influences.

In total there are about 13,500 feet of bank armor in Reach B3, which covers almost 30 percent of the bankline. Most of the armor is rock riprap, although there are over 3,000 feet of flow deflectors mapped in the reach, as well as over a mile of floodplain dikes.

Prior to 1950, 11,000 feet of side channels had been blocked in the reach, and since that time another 14,000 feet have been similarly blocked by small dikes. These ~4 miles of blocked channel are about equivalent in length to that of the main river. That said, as of 2001 there were still about 35,000 feet of active side channel in Reach B3.

Solid waste dumps were mapped on old side channels on the east floodplain areas at RM 361.5 and RM 360.6. There is one major headgate on the left bank of the river that feeds a heavily armored canal at RM 359.9.

Flow alterations and channel blockages have promoted the encroachment of riparian vegetation into old channel areas. Since 1950, almost 200 acres of riparian vegetation colonized previously un-vegetated side channels. Floodplain turnover rates have gone down since 1976 by about 2 acres per year, indicating slower rates of erosion.

Since 1950, predominantly agricultural land uses in Reach B3 have been converted to a mix of agriculture and urban/exurban development. About 1,000 acres of urban/exurban development has taken place since 1950. About 470 acres of ground continues to be flood irrigated in this area of east Billings. Approximately 16 percent of the Channel Migration Zone has become restricted due to physical features, all of which are bank armor installations designed to protect urban/industrial and agricultural land uses.

About 50 acres of Russian olive have been mapped in Reach B3. There are also fairly extensive mapped wetlands, with about 230 acres of total wetland area mapped, 95 acres of which are emergent wet meadows and marsh areas.

Reach B3 was sampled as part of the fisheries study. A total of 29 fish species were sampled in the reach, and none of those species have been identified by the Montana Natural Heritage Program as a Species of Concern (SOC).

Reach B3 was sampled as part of the avian study. The average species richness in this reach was 7.5, which indicates the average number of species observed during site visits to the reach in cottonwood habitats. The average species richness for sites evaluated is 8. One bird species identified by the Montana Natural Heritage Program as Potential Species of Concern (PSOC) was also found, the Plumbeous Vireo.

A hydrologic evaluation of flow depletions indicates that flow alterations over the last century have been substantial in this reach. The mean annual flood is estimated to have dropped from 23,900 cfs to 19,800 cfs, a drop of about 17 percent. The 2-year flood, which strongly influences overall channel form, has dropped from 44,500 cfs to 40,100 cfs, which is a reduction of 10 percent. Low flows have also been impacted; severe low flows described as 7Q10 (the lowest average 7-day flow anticipated every ten years) for summer months has dropped from an estimated 2,920 cfs to 2,010 cfs with human development, a reduction of 31 percent. More typical summer low flows, described as the summer 95% flow duration, have dropped from 3,836 cfs under unregulated conditions to 2,227 cfs under regulated conditions at the Billings gage, a reduction of 42 percent.

CEA-Related observations in Reach B3 include:

- Riparian encroachment with flow alterations
- Extensive armoring with CMZ encroachment

Recommended Practices (may include Yellowstone River Recommended Practices--YRRPs) for Reach B3 include:

- Side channel reactivation at RM 362.0, 360.5, 359.8 and RM 359.0
- Russian olive removal
- Solid waste dump removal RM 361.5 and RM 360.6
- Irrigation diversion structure management at RM 359.9.

PHYSICAL FEATURES MAP (2011)



HYDROLOGIC SUMMARY

Hydrologic data available for the Reach Narratives include data from representative gaging stations, modeling from the COE from the Big Horn river upstream, and modeling by the USGS for the Big Horn River to the Missouri River confluence. Gaging stations that best represent the watershed area within any reach are used to describe the flood history within the reach. Hydrology modeling results generated for all reaches provides unregulated and regulated flow values. Seasonal and annual flow duration data generated by the USGS are available for reaches C10 through D13.

Gage Representation (Gage-Based): Billings

Flood History

Year	Date	Flow on Date	Return Interval	Gage No	Downstream Gage	Upstream Gage
1943	Jun 21	61,200	10-25 yr		6309000	6214500
1996	Jun 12	61,900	10-25 yr		Miles City	Billings
1944	Jun 27	64,800	10-25 yr		1929-2015	1929-2015
1967	Jun 16	66,100	10-25 yr		Distance To (miles)	173.9
1975	Jul 7	67,600	10-25 yr			2.2
1974	Jun 19	69,500	25-50 yr			
2011	Jul 2	70,600	25-50 yr			
1918	Jun 15	78,100	50-100 yr			
1997	Jun 12	82,000	>100 yr			

Discharge

	1.01 Yr	2 Yr	5 Yr	10 Yr	50 Yr	100 Yr	500 Yr	7Q10 Summer	95% Sum. Duration
Unregulated	23,900	44,500	55,000	61,300	74,000	79,200	90,700	2,920	3,846
Regulated	19,800	40,100	50,900	57,500	71,000	76,600	89,100	2,010	2,227
% Change	-17.15%	-9.89%	-7.45%	-6.20%	-4.05%	-3.28%	-1.76%	-31.16%	-42.10%

AERIAL PHOTOGRAPHY

A variety of aerial photographic sources provide the basis for much of the Cumulative Effects Assessment analysis. The table below lists the air photos compiled for the reach and the associated discharge at the most representative USGS gaging station.

	Source	Acquisition Date	Type	Scale	Gage	Discharge
1950	USGS-EROS	14-May-51	B/W	1:28,400	6214500	13200
1976	USCOE	29-Sep-76	B/W	1:24,000	6214500	5630
1995	USGS DOQQ	23-Aug-96	B/W		6214500	4500
2001	NRCS	August 2-8, 2001	CIR	1:24,000	6214500	1700
2004	Merrick	15-May-04	Color	1:15,840	6214500	5960
2005	NAIP	07/08/2005	color	1-meter pixels	6214500	11400
2009	NAIP	7/5/2009	Color	1-meter pixels	6214500	23800
2011	USCOE	October 2012	color	1-ft pixel	6214500	3860
2011	NAIP	7/24/2011	Color	1-meter pixels	6214500	22800
2013	NAIP	06/15/2013	color	1-meter pixels	6214500	

PHYSICAL FEATURES

Several efforts to capture the types and extents of physical features in the corridor have been generated by the CEA study. The 2001 Physical Features Inventory was performed through helicopter/video Rapid Aerial Assessment by the NRCS (NRCS, 2001) and did not include Park County. This inventory includes point and linear features that represent bank armor, irrigation structures, transportation encroachments, and areas of accelerated erosion. Bank armor mapped in the 2001 inventory only reflects features on the active channel margin, and thus excludes off-channel features on historic side channels. Some floodplain restriction features such as dikes and levees in the 2001 Physical Features Inventory may extend well beyond the active channel. In 2013, the 2001 inventory was revised to include Park County. At that time, some attribute inconsistencies in the original data were addressed. This dataset was then updated to reflect conditions in the 2011 NAIP imagery.

For Stillwater, Yellowstone and Dawson Counties, a Physical Features Timeline was generated that includes additional mapping based on aerial photography and assigns approximate dates of feature construction based on observed presence/absence in historic imagery between the 1950s and 2005 (DTM and AGI, 2008). The Physical Features Timeline contains features that were not mapped in the 2001 inventory (e.g. bank armor abandoned in floodplain areas by 2001). As such the total bank armor extent in the 2005 data is commonly greater than that identified in 2001 or 2013.

Note: As the goal for each physical features mapping effort were different, with differing mapping extents, there will be discrepancies between total feature lengths (e.g. length of rock riprap) in each data set.

2001 and 2011 Physical Features Bankline Inventories

Feature Class	Feature Type	2001 Length (ft)	% of Bankline	2011 Length (ft)	% of Bankline	2001-2011 Change
Stream Stabilization						
	Rock RipRap	10,299	22.3%	10,047	21.7%	-252
	Flow Deflectors	731	1.6%	772	1.7%	41
	Concrete RipRap	592	1.3%	592	1.3%	0
	Between Flow Deflectors	2,338	5.1%	2,340	5.1%	1
	Feature Type Totals	13,960	30.2%	13,751	29.7%	-209
Floodplain Control						
	Transportation Encroachment	5,175	11.2%	5,175	11.2%	0
	Floodplain Dike/Levee	5,766	12.5%	5,766	12.5%	0
	Feature Type Totals	10,941	23.7%	10,941	23.7%	0
	Reach Totals	24,901	53.8%	24,692	53.4%	-209

Intent of Bank Protection: 2001

The 2001 bank protection features were assessed for the 'intent' of what they protect.

Feature Type	Irrigated	Non-Irrig.	Ag. Infrastr.	Road	Interstate	Railroad	Urban	Exurban
Concrete RipRap	0	98	0	0	0	0	0	886
Flow Deflectors/Between FDs	1,351	0	0	0	0	0	1,535	1,492
Rock RipRap	0	0	0	0	0	0	3,123	0
Totals	1,351	98	0	0	0	0	4,658	2,378

Bankline/Floodplain Inventory: Time Series


The Human Impacts Timeline assessed physical feature development through time for Yellowstone, Stillwater, and Dawson Counties.

Feature Class	Feature Type	Sum of Feature Length (ft)					
		1950	1976	1995	2001	2004	2005
Irrigation							
	Floodplain Dike/Levee	6,788	7,002	7,002	7,002	7,002	7,002
	Totals	6,788	7,002	7,002	7,002	7,002	7,002
Other							
	Floodplain Dike/Levee	7,446	7,446	7,446	7,446	7,446	7,446
	Totals	7,446	7,446	7,446	7,446	7,446	7,446
Other Off Channel							
	Floodplain Dike/Levee	0	2,866	6,494	6,494	6,494	6,494
	Floodplain Dike/Levee	155	7,025	9,010	9,010	9,010	9,010
	Totals	155	9,891	15,504	15,504	15,504	15,504
Stream Stabilization							

Rock RipRap	1,755	6,280	10,177	10,177	10,177	10,177
Flow Deflector	0	3,244	3,244	3,244	3,244	3,244
Concrete RipRap	0	0	592	592	592	592
Totals	1,755	9,524	14,012	14,012	14,012	14,012
Transportation Encroachment						
Railroad	5,149	5,149	5,149	5,149	5,149	5,149
Other	303	303	303	3,060	5,072	5,072
County Road	5,505	5,505	5,505	5,505	5,505	5,505
Totals	10,957	10,957	10,957	13,714	15,726	15,726

ICE JAMS

Ice jam data were obtained from the National Ice Jam Database maintained by the Ice Engineering Group at Army Corps of Engineers Cold Regions Research and Engineering Laboratory (<https://rsgis.crrel.usace.army.mil/icejam/>). From this database, Yellowstone River ice jams are summarized by reach in the Yellowstone River Historic Events Timeline (DTM and AGI, 2008b). The basic information for each ice jam is presented as a list of events. The graph represents the number of database entries for a reach. Note that a single jam event may have multiple entries.



GEOMORPHIC

The geomorphology data presented below consist of measured changes in Braiding Parameter since 1950 and blocked side channels. Braiding parameter is a measure of the total length of side channels relative to that of the main channel. The braiding parameter is calculated as the sum of anabranching and primary channel lengths divided by the primary channel length. Secondary channels within the bankfull margins are a function of flow stage and hence were not included in the braiding parameter calculation. If a reach has a braiding parameter of 3, then the total bankfull channel length is three times that of the main channel. The mean braiding parameter measured for all 88 reaches is 1.8.

Blocked side channels that were either plugged with a small dike or cutoff by larger features such as a levee or road prism were identified for the pre and post-1950s eras.

Additional geomorphic parameters are discussed in more detail in the study report and appendices.

Braiding (Bankfull)

	Primary Chan. Length (ft)	Anab. Ch. Length (ft)	Bankfull Braiding Parameter		% Change in Braiding
1950	22,668	41,147	2.82	1950 to 1976:	-21.52%
1976	23,157	28,007	2.21	1976 to 1995:	11.22%
1995	22,999	33,516	2.46	1995 to 2001:	2.60%
2001	23,124	35,173	2.52	1950 to 2001:	-10.45%
Change 1950 - 2001	456	-5,974	-0.29		

Length of Side Channels Blocked

Pre-1950s (ft)	11,002
Post-1950s (ft)	13,693

HYDRAULICS

Available hydraulic information includes county-based HEC-RAS modeling efforts by the Army Corps of Engineers with the exclusion of Park County. Floodplain modeling was performed for four conditions representing a developed and undeveloped floodplain, and unregulated and regulated flows for the 1.5, 2, 5, 10, 20, 50, 100, 200, and 500-year events. Park County has limited FEMA hydraulic modeling and was not included in the analysis.

The results of HEC-RAS modeling for the 5 and 100-year flood events were assessed to compare the extents of inundated area for the pristine (undeveloped floodplain, unregulated flows) and developed (developed floodplain, regulated flows) conditions. The data sets provided for each flow condition were unioned in the GIS to identify areas where the inundated extent differed. These areas of human-caused floodplain isolation due to either flow alterations or physical features such as levees. For the 100-year flood event, isolated areas greater than 5 acres were attributed with the interpreted reason for isolation (railroad, levee, etc.). The resulting values are presented as acres and percent of the pristine floodplain that has been isolated. The pristine floodplain is defined as the total floodplain footprint minus the area of the mapped 2001 bankfull channel (mapped islands were included in the floodplain area).

Floodplain Isolation

	100-Year		5-Year	
	Isolated Acres	% of Floodplain	Isolated Acres	% of Floodplain
Non-Structural (hydrology, geomorphic, etc.)	0	0.0%		
Agriculture (generally relates to field boundaries)	0	0.0%		
Agriculture (isolated by canal or large ditch)	0	0.0%		
Levee/Riprap (protecting agricultural lands)	0	0.0%		
Levee/Riprap (protecting urban, industrial, etc.)	0	0.0%		
Railroad	0	0.0%		
Abandoned Railroad	0	0.0%		
Transportation (Interstate and other roads)	0	0.0%		
Total Not Isolated (Ac)	1640		1489	
Total Floodplain Area (Ac)	1640		1644	
Total Isolated (Ac)	0	0.0%	155	14.1%

The 5-year floodplain is a good allegory for the extent of the riparian zone. Thus, irrigated areas within the 5-year floodplain tend to represent riparian zones that have been converted to agriculture and may result in additional bank protection to protect the agricultural production and irrigation infrastructure.

	Flood	Sprinkler	Pivot	Total
Irrigated Acres within the 5 Year Flooplain:	76	0	0	76

CHANNEL MIGRATION ZONE

A series of Channel Migration Maps were developed for the Yellowstone River from Gardiner to its mouth in McKenzie County, North Dakota (Thatcher, Swindell, and Boyd, 2009). These maps and their accompanying report can be accessed from the YRCDC Website. The channel migration zone (CMZ) developed for the Yellowstone River is defined as a composite area made up of the existing channel, the historic channel since 1950 (Historic Migration Zone, or HMZ), and an Erosion Buffer that encompasses areas prone to channel erosion over the next 100 years. Areas within this CMZ that have been isolated by constructed features such as armor or floodplain dikes are attributed as “Restricted Migration Areas” (RMA). Beyond the CMZ boundaries, outlying areas that pose risks of channel avulsion are identified as “Avulsion Potential Zones”.

Mean 50-Yr Migration Distance (ft)	Erosion Buffer (ft)	Total CMZ Acreage	Restricted CMZ Acreage	% Restricted Migration Area	Total AHZ Acreage	Restricted AHZ Acreage	% Restricted Avulsion Area
415	830	1,560	201	13%	64	64	100%

2011 Restricted Migration Area Summary

Note that these data reflect the observed conditions in the 2011 aerial photography (NAIP for Park and Sweet Grass Counties, COE for the rest of the river).

Reason for Restriction	Land Use Protected	RMA Acres	Percent of CMZ
RipRap/Flow Deflectors			
	Urban Industrial	105	6.4%
	Irrigated	129	7.9%
RipRap			
	Railroad	32	2.0%
	Totals	266	16.3%

Land Uses within the CMZ (Acres)

Flood Irrigation	Sprinkler Irrigation	Pivot Irrigation	Urban/ExUrban	Transportation
60.5	0.0	0.0	216.9	5.5

LAND USE

Land uses were mapped from aerial photography Gardiner to the confluence of the Missouri River in North Dakota for four time periods: 1950s, 1976, 2001, and 2011. Mapping was performed at approximately 1:6,000 to ensure consistent mapping across all data sets. Typically, if a feature could not be easily mapped at the target mapping scale, it was not separated out from the adjacent land use.

A four-tiered system was used to allow analysis at a variety of levels. Tier 1 breaks land use into Agricultural and Non-Agricultural uses. Tier two subdivided uses into productive Agricultural Land and Infrastructure for the Agricultural land, and Urban, Exurban and Transportation categories for the Non-Agricultural land. Tier three further breaks down land uses into more refined categories such as Irrigated or Non-Irrigated and Residential, Commercial, or Industrial. Finally, Tier 4 focuses primarily on the productive agricultural lands, identifying the type of irrigation (Pivot, Sprinkler or Flood).

Land Use Timeline - Tiers 2 and 3

Feature Class	Feature Type	Acres				% of Reach Area			
		1950	1976	2001	2011	1950	1976	2001	2011
Agricultural Infrastructure									
	Canal	0	0	0	0	0.0%	0.0%	0.0%	0.0%
	Agricultural Roads	0	0	0	0	0.0%	0.0%	0.0%	0.0%
	Other Infrastructure	50	67	71	51	1.3%	1.8%	1.9%	1.4%
	Totals	50	67	71	51	1.3%	1.8%	1.9%	1.4%
Agricultural Land									
	Non-Irrigated	2,297	1,703	1,387	1,297	60.5%	44.9%	36.5%	34.2%
	Irrigated	420	703	637	473	11.1%	18.5%	16.8%	12.4%
	Totals	2,717	2,406	2,025	1,770	71.6%	63.4%	53.3%	46.6%
Channel									
	Channel	870	906	837	853	22.9%	23.9%	22.1%	22.5%
	Totals	870	906	837	853	22.9%	23.9%	22.1%	22.5%
ExUrban									
	ExUrban Other	0	0	0	0	0.0%	0.0%	0.0%	0.0%
	ExUrban Undeveloped	0	0	0	0	0.0%	0.0%	0.0%	0.0%
	ExUrban Industrial	18	18	332	559	0.5%	0.5%	8.7%	14.7%
	ExUrban Commercial	0	0	0	0	0.0%	0.0%	0.0%	0.0%
	ExUrban Residential	3	14	39	57	0.1%	0.4%	1.0%	1.5%
	Totals	21	32	371	616	0.6%	0.8%	9.8%	16.2%
Transportation									
	Public Road	0	0	0	0	0.0%	0.0%	0.0%	0.0%
	Interstate	0	0	0	0	0.0%	0.0%	0.0%	0.0%
	Railroad	21	21	20	20	0.6%	0.6%	0.5%	0.5%
	Totals	21	21	20	20	0.6%	0.6%	0.5%	0.5%
Urban									
	Urban Other	0	0	40	27	0.0%	0.0%	1.1%	0.7%
	Urban Residential	0	96	171	182	0.0%	2.5%	4.5%	4.8%
	Urban Commercial	0	0	0	0	0.0%	0.0%	0.0%	0.0%
	Urban Undeveloped	0	12	0	13	0.0%	0.3%	0.0%	0.4%
	Urban Industrial	116	256	261	263	3.1%	6.8%	6.9%	6.9%
	Totals	116	365	473	485	3.1%	9.6%	12.5%	12.8%

Land Use Timeline - Tiers 3 and 4

Feature Class	Feature Type	Acres				% of Reach Area				Change Between Years (% of Agricultural Land)			
		1950	1976	2001	2011	1950	1976	2001	2011	'50-76	'76-01	'01-11	'50-11
Irrigated													
	Sprinkler	0	0	0	0	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
	Pivot	0	0	0	0	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
	Flood	420	703	637	473	15.5%	29.2%	31.5%	26.7%	13.7%	2.3%	-4.8%	11.2%
	Totals	420	703	637	473	15.5%	29.2%	31.5%	26.7%	13.7%	2.3%	-4.8%	11.2%

Non-Irrigated

Multi-Use	1,401	1,252	1,137	1,023	51.5%	52.0%	56.2%	57.8%	0.5%	4.1%	1.7%	6.3%
Hay/Pasture	896	451	250	274	33.0%	18.8%	12.4%	15.5%	-14.2%	-6.4%	3.1%	-17.5%
Totals	2,297	1,703	1,387	1,297	84.5%	70.8%	68.5%	73.3%	-13.7%	-2.3%	4.8%	-11.2%

RIPARIAN

Riparian mapping data are derived from the Yellowstone River Riparian Vegetation Mapping study (DTM/AGI 2008). This study coarsely mapped the riparian vegetation communities using 1950's, 1976-1977, and 2001 aerial imagery in a GIS environment. The polygons are digitized at a scale of approximately 1:7,500, with a minimum mapping unit of approximately 10 acres. The goal of the delineation was to capture areas of similar vegetation structure as they appeared on the aerial imagery, while maintaining a consistent scale.

The "Riparian Turnover" values quantify the total area within the active channel area that converted from either woody vegetation to open bar or water, or from open bar or water to woody vegetation. A comparison of these values allows some consideration of overall riparian encroachment into the river corridor from 1950 to 2001.

Riparian Mapping

Statistic	Shrub (Acres)			Closed Timber (Acres)			Open Timber (Acres)		
	1950	1976	2001	1950	1976	2001	1950	1976	2001
Min	4.4	0.5	1.0	1.6	1.4	1.2	0.7	2.6	1.7
Max	74.7	195.9	173.6	147.6	90.2	152.0	91.3	42.9	89.2
Average	29.4	13.8	22.3	29.9	20.3	32.7	20.9	17.8	36.0
Sum	205.9	385.2	356.1	448.2	507.7	523.3	292.9	106.5	179.9

Riparian Turnover

Conversion of riparian areas to channel, or from channel to riparian between the 1950's and 2001 data set.

Riparian to Channel (acres) 156.7

Channel to Riparian (acres) 214.1

Riparian Encroachment (acres) 57.3

Riparian Recruitment

Creation of riparian areas between 1950s and 2001.	1950s Channel Mapped as 2011 Riparian (Ac)	216.9
	1950s Floodplain Mapped as 2011 Channel (Ac)	138.0
	Total Recruitment (1950s to 2011)(Ac)	355.0

WETLANDS

Wetland areas were mapped to National Wetland Inventory standards by the Montana Natural Heritage Program. Palustrine wetlands within the mapped 100-year inundation boundary were extracted and summarized into four categories: Riverine (Unconsolidated Bottom - UB, Aquatic Bed - AB, and Unconsolidated Shore - US), Emergent - EM, Scrub-Shrub - SS, and Forested - FO.

	Riverine	Emergent	Scrub/Shrub	Forested	Total
Mapped Acres	95.8	94.9	40.5	0.0	231.2
Acres/Valley Mile	25.3	25.0	10.7	0.0	

RUSSIAN OLIVE

Russian olive is considered an invasive species and its presence in the Yellowstone River corridor is fairly recent. As such, its spread can be used as a general indicator of invasive plants within the corridor. It has the added benefit of being easily identified in multi-spectral aerial photography, making it possible to inventory large areas using remote techniques.

In 2011, Natural Resources Conservation Service (NRCS) in Bozeman, MT conducted an inventory of Russian olive locations in the Yellowstone River watershed. This study utilized the Feature Analyst extension within ArcGIS to interpret multi-spectral 2008 NAIP imagery for the presence of Russian olive. The resulting analysis was converted from raster format to a polygon ESRI shape file for distribution and further analysis within a GIS environment.

This work scope was tasked with integrating the resulting Russian olive inventory into the Yellowstone River Conservation Districts Council (YRDC) Cumulative Effects Assessment (CEA) GIS and associated reach-based database. Additionally, analysis of Russian olive within the corridor was conducted to characterize its distribution in throughout the corridor and its association with other corridor data sets.

	Floodplain Area (Ac)	% of Floodplain	Other Area (Ac)	Inside RMA (Ac)	Inside '50s Channel (Ac)	Inside 50s Island (Ac)
Russian Olive in Reach	49.76	4.14%	45.71	7.40	11.57	5.58

FISHERIES SUMMARY

Fisheries data available for the Reach Narratives include low-flow and high-flow habitat mapping of 2001 conditions for 406 miles of river, extending from the mouth upstream to a point approximately 8 miles upstream of Park City. Habitat mapping was performed remotely on the 2001 CIR aerial photography utilizing habitat classifications developed by Montana Fish, Wildlife, and Parks (DTM 2009). Historic habitat mapping using the 1950's imagery is limited to Reach B1 (high-flow) and D9 (low and high-flow).

Fisheries field sampling data have been provided by Ann Marie Reinhold (MSU). In this study, the Yellowstone River from Park City to Sidney was divided into five segments. Within each segment, fish were sampled in reaches modified by riprap ("treatment reaches") and relatively unmodified reaches ("control reaches"). Fish sampling was conducted during summer and autumn of 2009, 2010, and 2011. Boat electrofishing, trammel nets, mini-fyke nets and bag seines were used to collect data from river bends.

Fish presence data is only presented for those reaches that were sampled.

The Low Flow Habitat Mapping followed schema developed by Montana Fish Wildlife and Parks to identify key habitat units for certain aquatic species.

Fish Species Observed in Reach/Region

Species of Concern

Reach	Region	Reach	Region	Reach	Region	Reach	Region
<input type="checkbox"/>	<input checked="" type="checkbox"/> Bigmouth buffalo	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/> Flathead chub	<input type="checkbox"/>	<input type="checkbox"/> Northern redbelly dace	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/> Stonecat
<input type="checkbox"/>	<input checked="" type="checkbox"/> Black bullhead	<input type="checkbox"/>	<input type="checkbox"/> Freshwater drum	<input type="checkbox"/>	<input type="checkbox"/> Pallid sturgeon	<input type="checkbox"/>	<input type="checkbox"/> Sturgeon chub
<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/> Black crappie	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/> Goldeye	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/> Pumpkinseed	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/> Sucker species
<input type="checkbox"/>	<input type="checkbox"/> Blue sucker	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/> Green sunfish	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/> Rainbow trout	<input type="checkbox"/>	<input type="checkbox"/> Sunfish species
<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/> Bluegill	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/> Lake chub	<input type="checkbox"/>	<input checked="" type="checkbox"/> River carpsucker	<input type="checkbox"/>	<input type="checkbox"/> Walleye
<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/> Brook stickleback	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/> Largemouth bass	<input type="checkbox"/>	<input type="checkbox"/> Rock bass	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/> Western silvery minnow
<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/> Brown trout	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/> Longnose dace	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/> Sand shiner	<input type="checkbox"/>	<input type="checkbox"/> White bass
<input type="checkbox"/>	<input checked="" type="checkbox"/> Burbot	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/> Longnose sucker	<input type="checkbox"/>	<input checked="" type="checkbox"/> Sauger	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/> White crappie
<input type="checkbox"/>	<input type="checkbox"/> Catfish species	<input type="checkbox"/>	<input checked="" type="checkbox"/> Minnow species	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/> Shorthead redhorse	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/> White sucker
<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/> Channel catfish	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/> Mottled sculpin	<input type="checkbox"/>	<input type="checkbox"/> Shortnose gar	<input type="checkbox"/>	<input type="checkbox"/> Yellow bullhead
<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/> Common carp	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/> Mountain sucker	<input type="checkbox"/>	<input type="checkbox"/> Shovelnose sturgeon	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/> Yellow perch
<input type="checkbox"/>	<input type="checkbox"/> Creek chub	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/> Mountain whitefish	<input type="checkbox"/>	<input type="checkbox"/> Sicklefin chub		
<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/> Emerald shiner	<input type="checkbox"/>	<input type="checkbox"/> Northern pike	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/> Smallmouth bass		
<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/> Fathead minnow	<input type="checkbox"/>	<input type="checkbox"/> Northern plains killifish	<input type="checkbox"/>	<input type="checkbox"/> Smallmouth buffalo		

Low Flow Fisheries Habitat Mapping

2001 (Acres)

Habitat	Bankfull	Low Flow	% of Low Flow
Scour Pool	48.5	45.2	5.4%
Rip Rap Bottom	95.6	52.8	6.3%
Rip Rap Margin	28.3	13.1	1.6%
Secondary Channel	40.8	15.5	1.8%
Secondary Channel (Seasonal)	211.2	126.3	15.1%
Channel Crossover	116.1	47.6	5.7%
Point Bar		27.0	3.2%
Side Bar		44.3	5.3%
Mid-channel Bar		8.4	1.0%
Island	296.7	296.7	35.4%
Dry Channel		160.1	19.1%

AVIAN

Birds were sampled in 2006 and 2007 by Danielle Jones of Montana State University. Point count methods were used at 304 randomly chosen sites in 21 braided or anabranching reaches. Each site was visited multiple times within a season, and sites were visited in both years. Birds were sampled in grassland, shrubland, and cottonwood forest habitats. Additional bird data was collected by Amy Cilimburg of Montana Audubon in summer 2012. High priority areas for data collection were identified with the assistance of the YRDC Technical Advisory Committee. The Audubon methodology recorded data for a wider variety of bird species relative to the MSU study, including raptors and waterfowl.

Bird Species Observed in Reach/Region		Species of Concern	Potential Species of Concern
<input checked="" type="checkbox"/> <input checked="" type="checkbox"/> American Robin	<input type="checkbox"/> <input checked="" type="checkbox"/> Chipping Sparrow	<input type="checkbox"/> <input type="checkbox"/> Killdeer	<input checked="" type="checkbox"/> <input checked="" type="checkbox"/> Song Sparrow
<input checked="" type="checkbox"/> <input checked="" type="checkbox"/> American Crow	<input type="checkbox"/> <input checked="" type="checkbox"/> Clay-collared Sparrow	<input type="checkbox"/> <input type="checkbox"/> Lark Bunting	<input type="checkbox"/> <input checked="" type="checkbox"/> Spotted Sandpiper
<input checked="" type="checkbox"/> <input checked="" type="checkbox"/> American Goldfinch	<input type="checkbox"/> <input type="checkbox"/> Cliff Swallow	<input type="checkbox"/> <input checked="" type="checkbox"/> Lark Sparrow	<input checked="" type="checkbox"/> <input checked="" type="checkbox"/> Spotted Towhee
<input type="checkbox"/> <input type="checkbox"/> American Kestrel	<input checked="" type="checkbox"/> <input checked="" type="checkbox"/> Common Grackle	<input checked="" type="checkbox"/> <input checked="" type="checkbox"/> Lazuli Bunting	<input type="checkbox"/> <input type="checkbox"/> Sharp-shinned Hawk
<input checked="" type="checkbox"/> <input checked="" type="checkbox"/> American Redstart	<input type="checkbox"/> <input type="checkbox"/> Common Merganser	<input checked="" type="checkbox"/> <input checked="" type="checkbox"/> Least Flycatcher	<input type="checkbox"/> <input checked="" type="checkbox"/> Swainson's Thrush
<input type="checkbox"/> <input checked="" type="checkbox"/> Bald Eagle	<input type="checkbox"/> <input type="checkbox"/> Common Nighthawk	<input type="checkbox"/> <input checked="" type="checkbox"/> Mallard	<input type="checkbox"/> <input type="checkbox"/> Sandhill Crane
<input type="checkbox"/> <input type="checkbox"/> Baltimore Oriole	<input type="checkbox"/> <input checked="" type="checkbox"/> Common Raven	<input type="checkbox"/> <input type="checkbox"/> Mountain Bluebird	<input checked="" type="checkbox"/> <input checked="" type="checkbox"/> Tree Swallow
<input checked="" type="checkbox"/> <input checked="" type="checkbox"/> Barn Swallow	<input checked="" type="checkbox"/> <input checked="" type="checkbox"/> Common Yellowthroat	<input checked="" type="checkbox"/> <input checked="" type="checkbox"/> Mourning Dove	<input type="checkbox"/> <input type="checkbox"/> Turkey Vulture
<input type="checkbox"/> <input checked="" type="checkbox"/> Belted Kingfisher	<input type="checkbox"/> <input type="checkbox"/> Cooper's Hawk	<input checked="" type="checkbox"/> <input checked="" type="checkbox"/> Northern Flicker	<input type="checkbox"/> <input type="checkbox"/> Upland Sandpiper
<input type="checkbox"/> <input checked="" type="checkbox"/> Black-billed Cuckoo	<input type="checkbox"/> <input checked="" type="checkbox"/> Dickcissel	<input type="checkbox"/> <input type="checkbox"/> Orchard Oriole	<input type="checkbox"/> <input checked="" type="checkbox"/> Vesper Sparrow
<input checked="" type="checkbox"/> <input checked="" type="checkbox"/> Black-billed Magpie	<input checked="" type="checkbox"/> <input checked="" type="checkbox"/> Downy Woodpecker	<input type="checkbox"/> <input checked="" type="checkbox"/> Osprey	<input checked="" type="checkbox"/> <input checked="" type="checkbox"/> Violet-green Swallow
<input checked="" type="checkbox"/> <input checked="" type="checkbox"/> Black-capped Chickadee	<input type="checkbox"/> <input type="checkbox"/> Eastern Bluebird	<input type="checkbox"/> <input checked="" type="checkbox"/> Ovenbird	<input checked="" type="checkbox"/> <input checked="" type="checkbox"/> Warbling Vireo
<input type="checkbox"/> <input checked="" type="checkbox"/> Black-and-white Warbler	<input type="checkbox"/> <input checked="" type="checkbox"/> Eastern Kingbird	<input checked="" type="checkbox"/> <input checked="" type="checkbox"/> Plumbeous Vireo	<input checked="" type="checkbox"/> <input checked="" type="checkbox"/> Western Kingbird
<input checked="" type="checkbox"/> <input checked="" type="checkbox"/> Black-headed Grosbeak	<input type="checkbox"/> <input type="checkbox"/> Eurasian Collared-dove	<input type="checkbox"/> <input checked="" type="checkbox"/> Red-headed Woodpecker	<input type="checkbox"/> <input checked="" type="checkbox"/> Western Meadowlark
<input checked="" type="checkbox"/> <input checked="" type="checkbox"/> Blue Jay	<input checked="" type="checkbox"/> <input checked="" type="checkbox"/> European Starling	<input type="checkbox"/> <input type="checkbox"/> Red-naped Sapsucker	<input checked="" type="checkbox"/> <input checked="" type="checkbox"/> Western Wood-pewee
<input type="checkbox"/> <input type="checkbox"/> Bobolink	<input checked="" type="checkbox"/> <input checked="" type="checkbox"/> Field Sparrow	<input type="checkbox"/> <input type="checkbox"/> Red Crossbill	<input checked="" type="checkbox"/> <input checked="" type="checkbox"/> White-breasted Nuthatch
<input type="checkbox"/> <input checked="" type="checkbox"/> Brewer's Blackbird	<input type="checkbox"/> <input type="checkbox"/> Franklin's Gull	<input type="checkbox"/> <input checked="" type="checkbox"/> Ring-necked Pheasant	<input checked="" type="checkbox"/> <input checked="" type="checkbox"/> White-throated Swift
<input checked="" type="checkbox"/> <input checked="" type="checkbox"/> Brown-headed Cowbird	<input type="checkbox"/> <input type="checkbox"/> Grasshopper Sparrow	<input type="checkbox"/> <input checked="" type="checkbox"/> Red-tailed hawk	<input type="checkbox"/> <input type="checkbox"/> Wild Turkey
<input type="checkbox"/> <input type="checkbox"/> Brown Creeper	<input checked="" type="checkbox"/> <input checked="" type="checkbox"/> Gray Catbird	<input type="checkbox"/> <input checked="" type="checkbox"/> Rock Dove	<input type="checkbox"/> <input checked="" type="checkbox"/> Wood Duck
<input type="checkbox"/> <input checked="" type="checkbox"/> Brown Thrasher	<input type="checkbox"/> <input type="checkbox"/> Great Blue Heron	<input checked="" type="checkbox"/> <input checked="" type="checkbox"/> Red-winged Blackbird	<input type="checkbox"/> <input type="checkbox"/> Yellow-bellied Sapsucker
<input checked="" type="checkbox"/> <input checked="" type="checkbox"/> Bullock's Oriole	<input type="checkbox"/> <input type="checkbox"/> Great Horned Owl	<input checked="" type="checkbox"/> <input checked="" type="checkbox"/> Red-eyed Vireo	<input type="checkbox"/> <input type="checkbox"/> Yellow-billed Cuckoo
<input type="checkbox"/> <input checked="" type="checkbox"/> Canada Goose	<input checked="" type="checkbox"/> <input checked="" type="checkbox"/> Hairy Woodpecker	<input type="checkbox"/> <input type="checkbox"/> Red-breasted Grosbeak	<input checked="" type="checkbox"/> <input checked="" type="checkbox"/> Yellow-breasted Chat
<input checked="" type="checkbox"/> <input checked="" type="checkbox"/> Cedar Waxwing	<input type="checkbox"/> <input checked="" type="checkbox"/> House Finch	<input type="checkbox"/> <input type="checkbox"/> Say's Phoebe	<input type="checkbox"/> <input type="checkbox"/> Yellow-headed Blackbird
<input type="checkbox"/> <input checked="" type="checkbox"/> Chimney Swift	<input checked="" type="checkbox"/> <input checked="" type="checkbox"/> House Wren	<input checked="" type="checkbox"/> <input checked="" type="checkbox"/> Savannah Sparrow	<input checked="" type="checkbox"/> <input checked="" type="checkbox"/> Yellow Warbler

CULTURAL INVENTORY SUMMARY

The Yellowstone River Cultural Inventory - 2006 documents the variety and intensity of different perspectives and values held by people who share the Yellowstone River. Between May and November of 2006, a total of 313 individuals participated in the study. They represented agricultural, civic, recreational, or residential interest groups. Also, individuals from the Crow and the Northern Cheyenne tribes were included.

There are three particular goals associated with the investigation. The first goal is to document how the people of the Yellowstone River describe the physical character of the river and how they think the physical processes, such as floods and erosion, should be managed. Within this goal, efforts have been made to document participants' views regarding the many different bank stabilization techniques employed by landowners. The second goal is to document the degree to which the riparian zone associated with the river is recognized and valued by the participants. The third goal is to document concerns regarding the management of the river's resources. Special attention is given to the ways in which residents from diverse geographical settings and diverse interest groups view river management and uses. The results illustrate the commonalities of thought and the complexities of concerns expressed by those who share the resources of the Yellowstone River.

Summary of Cultural Views in Region B

The study segment Big Horn to Laurel includes data from the people of one large county, Yellowstone County. Three themes dominate conversations with the four interest groups. One theme focuses on the evolving communities of Yellowstone County, most of which are influenced by the economic success and sheer growth of Billings. The second theme focuses on the evolving relationships that the people have with the river. While traditional agricultural activities continue in the county, many people discuss notions related to urban and residential experiences and how the river becomes an asset that improves one's quality of life as an urban dweller. The third theme involves a complex tangle of pressures and demands that require managerial strategies capable of dealing with a future that has arrived.