

County	Rosebud	Upstream River Mile	243.1
Classification	PCM: Partially confined meandering	Downstream River Mile	236.3
General Location	Forsyth	Length	6.80 mi (10.94 km)
General Comments	Forsyth		

Narrative Summary

Reach C10 is 6.8 miles long and is located at Forsyth. It is a Partially Confined Meandering reach type, as the river flows within a primary meandering thread that is partially confined by the northern bluff line at the Forsyth Bridge.

There is approximately three miles of rock riprap in the reach, 500 feet of which was built since 2001. About a mile of armor is protecting the active rail line on the south side of the river, and another 3,700 feet are protecting the city of Forsyth. Just below Cartersville Dam, a ~330 foot-long stretch of bank armor was flanked sometime between 2001 and 2011. The river has since migrated to the south about 50 feet past the abandoned armor. As of 2011 there were 1,600 feet of flow deflectors mapped in the reach. About 22 percent of the total bankline is armored by either rock riprap or flow deflectors. There is also about a mile of floodplain dikes/levees in the reach, which are located on the south bank at Forsyth.

Cartersville Dam is located at RM 238.5 in the town of Forsyth. This diversion dam was constructed in the early 1930's and consists of a rock rubble riprap core that is capped by concrete. The structure is 800 feet long, spanning the width of the Yellowstone River. The river flows within a single thread at the structure, flowing along the northern bluff line of the Yellowstone River valley. Because of its impacts on the Yellowstone River fishery, efforts have begun to develop suitable alternatives and bypass designs to promote fish passage at Cartersville.

About 20 percent of the total 100-year floodplain has become isolated due to human development. The isolation is due to a combination of floodplain dikes that protect the city of Forsyth and the active railroad. The 5-year floodplain is even more affected; 50 percent of the historic 5-year floodplain is no longer inundated at that frequency. Most of the isolated 5-year floodplain area is occupied by flood irrigated fields north of the river, and by urban development in Forsyth. At RM 238 the river is migrating northward, and has reached the toe of the abandoned Milwaukee Rail Line embankment. Migration through this grade will increase floodplain access on the north side of the river downstream of Cartersville Dam. As this is an urban reach, strategic floodplain reconnection in this area could be beneficial.

One ice jam was reported in Reach C10 in February of 1996. No damages were reported.

Land use is dominated by agriculture (~4,700 acres), with 280 acres of pivot irrigation development since 1950. There are about 850 acres of urban/exurban development in the reach. About 4 percent of the CMZ is restricted by physical features, and most of that area is in town.

There are 250 acres of Russian olive in the reach, most of which is dispersed in riparian areas. Russian olive densities are especially high downstream of Cartersville Diversion dam on the south bank of the river near the water treatment plant.

A hydrologic evaluation of flow depletions indicates that flow alterations over the last century have been major in this reach. The 2-year flood, which strongly influences overall channel form, has dropped by 24 percent. Low flows have also been impacted; severe low flows described as 7Q10 (the lowest average 7-day flow anticipated every ten years) for summer months has dropped from an estimated 4,730 cfs to 3,020 cfs with human development, a reduction of 36 percent. More typical summer low flows, described as the summer 95% flow duration, have dropped from 6,150 cfs under unregulated conditions to 3,320 cfs under regulated conditions, a reduction of 46 percent.


CEA-Related observations in Reach C10 include:

- Floodplain isolation due to urban/exurban development.
- Extensive Russian olive colonization in urbanized reach

Recommended Practices (may include Yellowstone River Recommended Practices--YRRPs) for Reach C10 include:

- Floodplain reconnection at RM 238L behind abandoned Milwaukee rail line.
- Diversion structure management at Cartersville Dam
- Watercraft passage at Cartersville Dam
- Fish Passage at Cartersville Dam
- Flanked bank armor removal at RM 238.4R
- Russian olive removal

PHYSICAL FEATURES MAP (2011)



HYDROLOGIC SUMMARY

Hydrologic data available for the Reach Narratives include data from representative gaging stations, modeling from the COE from the Big Horn river upstream, and modeling by the USGS for the Big Horn River to the Missouri River confluence. Gaging stations that best represent the watershed area within any reach are used to describe the flood history within the reach. Hydrology modeling results generated for all reaches provides unregulated and regulated flow values. Seasonal and annual flow duration data generated by the USGS are available for reaches C10 through D13.

Gage Representation (Gage-Based): Miles City

Flood History

Year	Date	Flow on Date	Return Interval	Gage No	Downstream Gage	Upstream Gage
1974	Jun 22	75,400	10-25 yr	6309000	6214500	
1997	Jun 15	83,300	10-25 yr	Miles City	Billings	
1943	Jun 26	83,700	10-25 yr	1929-2015	1929-2015	
2011	May 24	85,400	10-25 yr	Distance To (miles)	52.3	121.3
1944	Jun 19	96,300	50-100 yr			
1978	May 22	102,000	50-100 yr			

Discharge

	1.01 Yr	2 Yr	5 Yr	10 Yr	50 Yr	100 Yr	500 Yr	7Q10 Summer	95% Sum. Duration
Unregulated		61,300	77,300	87,900	111,000	121,000	145,000	4,730	6,150
Regulated		46,900	61,300	70,700	91,600	101,000	122,000	3,020	3,320
% Change		-23.49%	-20.70%	-19.57%	-17.48%	-16.53%	-15.86%	-36.15%	-46.02%

Flow Duration

Streamflow, in ft³/s, which was equaled or exceeded for indicated percent of time

Note that these statistics are only available from Reach C10 downstream. See the USGS report for detailed information.

Season		5%	50%	95%
Spring	Unregulated	60,000	22,400	5,930
	Regulated	46,500	13,600	4,330
	% Change	-23%	-39%	-27%
Summer	Unregulated	42,100	13,200	6,150
	Regulated	32,200	8,230	3,320
	% Change	-24%	-38%	-46%
Fall	Unregulated	9,030	5,460	2,280
	Regulated	10,400	6,800	3,590
	% Change	15%	25%	57%
Winter	Unregulated	11,400	4,850	1,990
	Regulated	12,000	5,940	3,230
	% Change	5%	22%	62%
Annual	Unregulated	44,900	7,770	2,760
	Regulated	33,800	7,280	3,580
	% Change	-25%	-6%	30%

AERIAL PHOTOGRAPHY

A variety of aerial photographic sources provide the basis for much of the Cumulative Effects Assessment analysis. The table below lists the air photos compiled for the reach and the associated discharge at the most representative USGS gaging station.

	Source	Acquisition Date	Type	Scale	Gage	Discharge
1950	USGS-EROS	26-Aug-49	B/W	1:14,800	6309000	3620
1976	USCOE	29-Sep-76	B/W	1:24,000	6309000	9520
1995	USGS DOQQ	11-Aug-96	B/W		6295000	7650
2001	NRCS	August 2-8, 2001	CIR	1:24,000	6295000	3500
2005	NAIP	07/12/2005	color	1-meter pixels	6309000	17500
2007	Woolpert	10/15/2007 - 11/2/0007	Color			
2009	NAIP	8/11/2009	Color	1-meter pixels	6309000	12900
2011	USCOE	October 2012	color	1-ft pixel	6309000	8100
2011	NAIP	7/15/2011	Color	1-meter pixels	6309000	58000
2013	NAIP	07/21/2013	color	1-meter pixels	6309000	

PHYSICAL FEATURES

Several efforts to capture the types and extents of physical features in the corridor have been generated by the CEA study. The 2001 Physical Features Inventory was performed through helicopter/video Rapid Aerial Assessment by the NRCS (NRCS, 2001) and did not include Park County. This inventory includes point and linear features that represent bank armor, irrigation structures, transportation encroachments, and areas of accelerated erosion. Bank armor mapped in the 2001 inventory only reflects features on the active channel margin, and thus excludes off-channel features on historic side channels. Some floodplain restriction features such as dikes and levees in the 2001 Physical Features Inventory may extend well beyond the active channel. In 2013, the 2001 inventory was revised to include Park County. At that time, some attribute inconsistencies in the original data were addressed. This dataset was then updated to reflect conditions in the 2011 NAIP imagery.

For Stillwater, Yellowstone and Dawson Counties, a Physical Features Timeline was generated that includes additional mapping based on aerial photography and assigns approximate dates of feature construction based on observed presence/absence in historic imagery between the 1950s and 2005 (DTM and AGI, 2008). The Physical Features Timeline contains features that were not mapped in the 2001 inventory (e.g. bank armor abandoned in floodplain areas by 2001). As such the total bank armor extent in the 2005 data is commonly greater than that identified in 2001 or 2013.

Note: As the goal for each physical features mapping effort were different, with differing mapping extents, there will be discrepancies between total feature lengths (e.g. length of rock riprap) in each data set.

2001 and 2011 Physical Features Bankline Inventories

Feature Class	Feature Type	2001 Length (ft)	% of Bankline	2011 Length (ft)	% of Bankline	2001-2011 Change
Stream Stabilization						
	Rock RipRap	13,814	19.2%	14,306	19.8%	493
	Flow Deflectors	607	0.8%	345	0.5%	-262
	Between Flow Deflectors	1,302	1.8%	1,302	1.8%	0
	Feature Type Totals	15,723	21.8%	15,954	22.1%	231
Floodplain Control						
	Floodplain Dike/Levee	4,861	6.7%	4,071	5.6%	-790
	Feature Type Totals	4,861	6.7%	4,071	5.6%	-790
	Reach Totals	20,584	28.6%	20,025	27.8%	-559


Intent of Bank Protection: 2001

The 2001 bank protection features were assessed for the 'intent' of what they protect.

Feature Type	Irrigated	Non-Irrig.	Ag. Infrastr.	Road	Interstate	Railroad	Urban	Exurban
Flow Deflectors/Between FDs	1,725	0	0	0	0	0	0	0
Rock RipRap	0	0	0	722	0	5,054	3,720	0
Totals	1,725	0	0	722	0	5,054	3,720	0

ICE JAMS

Ice jam data were obtained from the National Ice Jam Database maintained by the Ice Engineering Group at Army Corps of Engineers Cold Regions Research and Engineering Laboratory (<https://rsgis.crrel.usace.army.mil/icejam/>). From this database, Yellowstone River ice jams are summarized by reach in the Yellowstone River Historic Events Timeline (DTM and AGI, 2008b). The basic information for each ice jam is presented as a list of events. The graph represents the number of database entries for a reach. Note that a single jam event may have multiple entries.



Jam Date	Jam Type	River Mile	Damages
2/7/1996	Break-up	239	?

GEOMORPHIC

The geomorphology data presented below consist of measured changes in Braiding Parameter since 1950 and blocked side channels. Braiding parameter is a measure of the total length of side channels relative to that of the main channel. The braiding parameter is calculated as the sum of anabranching and primary channel lengths divided by the primary channel length. Secondary channels within the bankfull margins are a function of flow stage and hence were not included in the braiding parameter calculation. If a reach has a braiding parameter of 3, then the total bankfull channel length is three times that of the main channel. The mean braiding parameter measured for all 88 reaches is 1.8.

Blocked side channels that were either plugged with a small dike or cutoff by larger features such as a levee or road prism were identified for the pre and post-1950s eras.

Additional geomorphic parameters are discussed in more detail in the study report and appendices.

Braiding (Bankfull)

	Primary Chan. Length (ft)	Anab. Ch. Length (ft)	Bankfull Braiding Parameter	% Change in Braiding
1950	37,786	9,048	1.24	1950 to 1976: 3.26%
1976	35,535	9,945	1.28	1976 to 1995: 6.47%
1995	36,024	13,064	1.36	1995 to 2001: 5.39%
2001	36,044	15,719	1.44	1950 to 2001: 15.87%
Change 1950 - 2001	-1,742	6,671	0.20	

Length of Side Channels Blocked	Pre-1950s (ft)	Post-1950s (ft)
	0	0

HYDRAULICS

Available hydraulic information includes county-based HEC-RAS modeling efforts by the Army Corps of Engineers with the exclusion of Park County. Floodplain modeling was performed for four conditions representing a developed and undeveloped floodplain, and unregulated and regulated flows for the 1.5, 2, 5, 10, 20, 50, 100, 200, and 500-year events. Park County has limited FEMA hydraulic modeling and was not included in the analysis.

The results of HEC-RAS modeling for the 5 and 100-year flood events were assessed to compare the extents of inundated area for the pristine (undeveloped floodplain, unregulated flows) and developed (developed floodplain, regulated flows) conditions. The data sets provided for each flow condition were unioned in the GIS to identify areas where the inundated extent differed. These areas of human-caused floodplain isolation due to either flow alterations or physical features such as levees. For the 100-year flood event, isolated areas greater than 5 acres were attributed with the interpreted reason for isolation (railroad, levee, etc.). The resulting values are presented as acres and percent of the pristine floodplain that has been isolated. The pristine floodplain is defined as the total floodplain footprint minus the area of the mapped 2001 bankfull channel (mapped islands were included in the floodplain area).

Floodplain Isolation

	100-Year		5-Year	
	Isolated Acres	% of Floodplain	Isolated Acres	% of Floodplain
Non-Structural (hydrology, geomorphic, etc.)	44	1.4%		
Agriculture (generally relates to field boundaries)	0	0.0%		
Agriculture (isolated by canal or large ditch)	0	0.0%		
Levee/Riprap (protecting agricultural lands)	0	0.0%		
Levee/Riprap (protecting urban, industrial, etc.)	338	10.8%		
Railroad	223	7.1%		
Abandoned Railroad	16	0.5%		
Transportation (Interstate and other roads)	15	0.5%		
Total Not Isolated (Ac)	2507		1753	
Total Floodplain Area (Ac)	3143		2872	
Total Isolated (Ac)	636	20.2%	1119	49.9%

The 5-year floodplain is a good allegory for the extent of the riparian zone. Thus, irrigated areas within the 5-year floodplain tend to represent riparian zones that have been converted to agriculture and may result in additional bank protection to protect the agricultural production and irrigation infrastructure.

	Flood	Sprinkler	Pivot	Total
Irrigated Acres within the 5 Year Flooplain:	29	0	21	50

CHANNEL MIGRATION ZONE

A series of Channel Migration Maps were developed for the Yellowstone River from Gardiner to its mouth in McKenzie County, North Dakota (Thatcher, Swindell, and Boyd, 2009). These maps and their accompanying report can be accessed from the YRCDC Website. The channel migration zone (CMZ) developed for the Yellowstone River is defined as a composite area made up of the existing channel, the historic channel since 1950 (Historic Migration Zone, or HMZ), and an Erosion Buffer that encompasses areas prone to channel erosion over the next 100 years. Areas within this CMZ that have been isolated by constructed features such as armor or floodplain dikes are attributed as "Restricted Migration Areas" (RMA). Beyond the CMZ boundaries, outlying areas that pose risks of channel avulsion are identified as "Avulsion Potential Zones".

Mean 50-Yr Migration Distance (ft)	Erosion Buffer (ft)	Total CMZ Acreage	Restricted CMZ Acreage	% Restricted Migration Area	Total AHZ Acreage	Restricted AHZ Acreage	% Restricted Avulsion Area
210	420	1,344	67	5%	501	0	0%

2011 Restricted Migration Area Summary

Note that these data reflect the observed conditions in the 2011 aerial photography (NAIP for Park and Sweet Grass Counties, COE for the rest of the river).

Reason for Restriction	Land Use Protected	RMA Acres	Percent of CMZ
RipRap/Flow Deflectors	Irrigated	19	1.0%
RipRap	Urban Residential	11	0.6%
Dike/Levee	Urban Residential	43	2.3%
Totals		73	3.9%

Land Uses within the CMZ (Acres)

Flood Irrigation	Sprinkler Irrigation	Pivot Irrigation	Urban/ExUrban	Transportation
39.4	0.0	2.5	70.8	1.6

LAND USE

Land uses were mapped from aerial photography Gardiner to the confluence of the Missouri River in North Dakota for four time periods: 1950s, 1976, 2001, and 2011. Mapping was performed at approximately 1:6,000 to ensure consistent mapping across all data sets. Typically, if a feature could not be easily mapped at the target mapping scale, it was not separated out from the adjacent land use.

A four-tiered system was used to allow analysis at a variety of levels. Tier 1 breaks land use into Agricultural and Non-Agricultural uses. Tier two subdivided uses into productive Agricultural Land and Infrastructure for the Agricultural land, and Urban, Exurban and Transportation categories for the Non-Agricultural land. Tier three further breaks down land uses into more refined categories such as Irrigated or Non-Irrigated and Residential, Commercial, or Industrial. Finally, Tier 4 focuses primarily on the productive agricultural lands, identifying the type of irrigation (Pivot, Sprinkler or Flood).

Land Use Timeline - Tiers 2 and 3

Feature Class	Feature Type	Acres				% of Reach Area			
		1950	1976	2001	2011	1950	1976	2001	2011
Agricultural Infrastructure									
	Canal	0	0	0	0	0.0%	0.0%	0.0%	0.0%
	Agricultural Roads	0	0	0	0	0.0%	0.0%	0.0%	0.0%
	Other Infrastructure	29	72	101	104	0.4%	1.1%	1.5%	1.5%
	Totals	29	72	101	104	0.4%	1.1%	1.5%	1.5%
Agricultural Land									
	Non-Irrigated	4,488	3,771	3,602	3,565	67.0%	56.3%	53.8%	53.2%
	Irrigated	904	1,138	1,166	1,152	13.5%	17.0%	17.4%	17.2%
	Totals	5,392	4,909	4,768	4,717	80.5%	73.3%	71.2%	70.4%
Channel									
	Channel	684	736	706	758	10.2%	11.0%	10.5%	11.3%
	Totals	684	736	706	758	10.2%	11.0%	10.5%	11.3%
ExUrban									
	ExUrban Other	0	26	26	26	0.0%	0.4%	0.4%	0.4%
	ExUrban Undeveloped	0	0	21	0	0.0%	0.0%	0.3%	0.0%
	ExUrban Industrial	0	18	18	18	0.0%	0.3%	0.3%	0.3%
	ExUrban Commercial	0	0	0	0	0.0%	0.0%	0.0%	0.0%
	ExUrban Residential	0	1	76	97	0.0%	0.0%	1.1%	1.5%
	Totals	0	45	142	142	0.0%	0.7%	2.1%	2.1%
Transportation									
	Public Road	36	56	57	57	0.5%	0.8%	0.9%	0.9%
	Interstate	0	153	153	153	0.0%	2.3%	2.3%	2.3%
	Railroad	72	72	37	37	1.1%	1.1%	0.6%	0.6%
	Totals	107	281	248	248	1.6%	4.2%	3.7%	3.7%
Urban									
	Urban Other	102	70	106	102	1.5%	1.0%	1.6%	1.5%
	Urban Residential	270	365	390	390	4.0%	5.5%	5.8%	5.8%
	Urban Commercial	41	80	97	97	0.6%	1.2%	1.4%	1.4%
	Urban Undeveloped	66	44	41	41	1.0%	0.7%	0.6%	0.6%
	Urban Industrial	4	93	97	97	0.1%	1.4%	1.4%	1.5%
	Totals	484	652	732	728	7.2%	9.7%	10.9%	10.9%

Land Use Timeline - Tiers 3 and 4

Feature Class	Feature Type	Acres				% of Reach Area				Change Between Years (% of Agricultural Land)			
		1950	1976	2001	2011	1950	1976	2001	2011	'50-76	'76-01	'01-11	'50-11
Irrigated													
	Sprinkler	0	0	0	0	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
	Pivot	0	206	263	278	0.0%	4.2%	5.5%	5.9%	4.2%	1.3%	0.4%	5.9%
	Flood	904	932	904	874	16.8%	19.0%	18.9%	18.5%	2.2%	0.0%	-0.4%	1.8%
	Totals	904	1,138	1,166	1,152	16.8%	23.2%	24.5%	24.4%	6.4%	1.3%	0.0%	7.7%

Non-Irrigated

Multi-Use	4,015	3,577	3,585	3,557	74.4%	72.9%	75.2%	75.4%	-1.6%	2.3%	0.2%	1.0%
Hay/Pasture	474	194	17	8	8.8%	4.0%	0.4%	0.2%	-4.8%	-3.6%	-0.2%	-8.6%
Totals	4,488	3,771	3,602	3,565	83.2%	76.8%	75.5%	75.6%	-6.4%	-1.3%	0.0%	-7.7%

RIPARIAN

Riparian mapping data are derived from the Yellowstone River Riparian Vegetation Mapping study (DTM/AGI 2008). This study coarsely mapped the riparian vegetation communities using 1950's, 1976-1977, and 2001 aerial imagery in a GIS environment. The polygons are digitized at a scale of approximately 1:7,500, with a minimum mapping unit of approximately 10 acres. The goal of the delineation was to capture areas of similar vegetation structure as they appeared on the aerial imagery, while maintaining a consistent scale.

The "Riparian Turnover" values quantify the total area within the active channel area that converted from either woody vegetation to open bar or water, or from open bar or water to woody vegetation. A comparison of these values allows some consideration of overall riparian encroachment into the river corridor from 1950 to 2001.

Riparian Mapping

Statistic	Shrub (Acres)			Closed Timber (Acres)			Open Timber (Acres)		
	1950	1976	2001	1950	1976	2001	1950	1976	2001
Min	0.4	1.9	0.9	0.9	1.7	2.4	5.6	3.6	18.3
Max	294.3	241.2	171.7	241.2	281.1	163.9	232.9	115.9	116.4
Average	36.5	33.0	27.6	49.1	58.3	33.1	54.4	29.7	76.1
Sum	474.5	296.9	386.5	736.7	815.9	694.5	435.0	267.4	380.3

Riparian Turnover

Conversion of riparian areas to channel, or from channel to riparian between the 1950's and 2001 data set.

Riparian to Channel (acres) 87.1

Channel to Riparian (acres) 119.1

Riparian Encroachment (acres) 32.0

Riparian Recruitment

Creation of riparian areas between 1950s and 2001.	1950s Channel Mapped as 2011 Riparian (Ac)	128.0
	1950s Floodplain Mapped as 2011 Channel (Ac)	12.3
	Total Recruitment (1950s to 2011)(Ac)	140.3

WETLANDS

Wetland areas were mapped to National Wetland Inventory standards by the Montana Natural Heritage Program. Palustrine wetlands within the mapped 100-year inundation boundary were extracted and summarized into four categories: Riverine (Unconsolidated Bottom - UB, Aquatic Bed - AB, and Unconsolidated Shore - US), Emergent - EM, Scrub-Shrub - SS, and Forested - FO.

	Riverine	Emergent	Scrub/Shrub	Forested	Total
Mapped Acres	11.6	89.6	30.1	0.0	131.2
Acres/Valley Mile	1.9	14.8	5.0	0.0	

RUSSIAN OLIVE

Russian olive is considered an invasive species and its presence in the Yellowstone River corridor is fairly recent. As such, its spread can be used as a general indicator of invasive plants within the corridor. It has the added benefit of being easily identified in multi-spectral aerial photography, making it possible to inventory large areas using remote techniques.

In 2011, Natural Resources Conservation Service (NRCS) in Bozeman, MT conducted an inventory of Russian olive locations in the Yellowstone River watershed. This study utilized the Feature Analyst extension within ArcGIS to interpret multi-spectral 2008 NAIP imagery for the presence of Russian olive. The resulting analysis was converted from raster format to a polygon ESRI shape file for distribution and further analysis within a GIS environment.

This work scope was tasked with integrating the resulting Russian olive inventory into the Yellowstone River Conservation Districts Council (YRDC) Cumulative Effects Assessment (CEA) GIS and associated reach-based database. Additionally, analysis of Russian olive within the corridor was conducted to characterize its distribution in throughout the corridor and its association with other corridor data sets.

	Floodplain Area (Ac)	% of Floodplain	Other Area (Ac)	Inside RMA (Ac)	Inside '50s Channel (Ac)	Inside 50s Island (Ac)
Russian Olive in Reach	250.55	5.68%	6.77	2.27	15.11	1.46

FISHERIES SUMMARY

Fisheries data available for the Reach Narratives include low-flow and high-flow habitat mapping of 2001 conditions for 406 miles of river, extending from the mouth upstream to a point approximately 8 miles upstream of Park City. Habitat mapping was performed remotely on the 2001 CIR aerial photography utilizing habitat classifications developed by Montana Fish, Wildlife, and Parks (DTM 2009). Historic habitat mapping using the 1950's imagery is limited to Reach B1 (high-flow) and D9 (low and high-flow).

Fisheries field sampling data have been provided by Ann Marie Reinhold (MSU). In this study, the Yellowstone River from Park City to Sidney was divided into five segments. Within each segment, fish were sampled in reaches modified by riprap ("treatment reaches") and relatively unmodified reaches ("control reaches"). Fish sampling was conducted during summer and autumn of 2009, 2010, and 2011. Boat electrofishing, trammel nets, mini-fyke nets and bag seines were used to collect data from river bends.

Fish presence data is only presented for those reaches that were sampled.

The Low Flow Habitat Mapping followed schema developed by Montana Fish Wildlife and Parks to identify key habitat units for certain aquatic species.

Low Flow Fisheries Habitat Mapping

Habitat	2001 (Acres)		
	Bankfull	Low Flow	% of Low Flow
Scour Pool	180.9	105.4	14.9%
Rip Rap Bottom	31.5	21.8	3.1%
Rip Rap Margin	122.2	100.7	14.3%
Secondary Channel	28.5	22.9	3.2%
Secondary Channel (Seasonal)	71.5	41.6	5.9%
Channel Crossover	110.0	102.7	14.5%
Point Bar		55.3	7.8%
Side Bar		14.8	2.1%
Mid-channel Bar		28.2	4.0%
Island	76.1	76.1	10.8%
Dry Channel		51.4	7.3%
Dam Influenced	85.2	85.1	12.1%

AVIAN

Birds were sampled in 2006 and 2007 by Danielle Jones of Montana State University. Point count methods were used at 304 randomly chosen sites in 21 braided or anabranching reaches. Each site was visited multiple times within a season, and sites were visited in both years. Birds were sampled in grassland, shrubland, and cottonwood forest habitats. Additional bird data was collected by Amy Cilimburg of Montana Audubon in summer 2012. High priority areas for data collection were identified with the assistance of the YRCDC Technical Advisory Committee. The Audubon methodology recorded data for a wider variety of bird species relative to the MSU study, including raptors and waterfowl.

CULTURAL INVENTORY SUMMARY

The Yellowstone River Cultural Inventory - 2006 documents the variety and intensity of different perspectives and values held by people who share the Yellowstone River. Between May and November of 2006, a total of 313 individuals participated in the study. They represented agricultural, civic, recreational, or residential interest groups. Also, individuals from the Crow and the Northern Cheyenne tribes were included.

There are three particular goals associated with the investigation. The first goal is to document how the people of the Yellowstone River describe the physical character of the river and how they think the physical processes, such as floods and erosion, should be managed. Within this goal, efforts have been made to document participants' views regarding the many different bank stabilization techniques employed by landowners. The second goal is to document the degree to which the riparian zone associated with the river is recognized and valued by the participants. The third goal is to document concerns regarding the management of the river's resources. Special attention is given to the ways in which residents from diverse geographical settings and diverse interest groups view river management and uses. The results illustrate the commonalities of thought and the complexities of concerns expressed by those who share the resources of the Yellowstone River.

Summary of Cultural Views in Region C

In the study segment, Powder River to Big Horn River, three conversations emerged across the four interest groups. The first conversation focuses on the "familiar way of life." The conversation exposes a local identity that is tied to agriculture and to traditional forms of recreation, such as hunting and fishing. When asked if the familiar management practices are sufficient in terms of sharing the river's resources, some locals express concerns. The second conversation explicitly acknowledges that the demand for recreational access to the river's resources is in its infancy in terms of representing a problem. The third conversation focuses on controlling the river with rip-rap and dikes.