

<b>County</b>	Dawson	<b>Upstream River Mile</b>	89
<b>Classification</b>	PCA: Partially confined anabranching	<b>Downstream River Mile</b>	81.4
<b>General Location</b>	Downstream of Glendive	<b>Length</b>	7.60 mi (12.23 km)

## General Comments

### Narrative Summary

Reach D7 is located just downstream of Glendive. It is 7.6 miles long and is a Partially Confined Anabranching (PCA) reach type, including some valley wall influence as well as numerous forested islands. These reach types tend to be relatively dynamic with high rates of channel change through time. The Stipek Fishing Access Site is located in the middle portion of the reach.

No bank armor has been mapped in Reach D7, and no side channels have been blocked by dikes. About two miles of transportation encroachment by the railroad was mapped in Reach D7, all of which was in place by 1950.

Similar to many reaches in the Lower Yellowstone Valley, the river channel in Reach D7 has gotten smaller since 1950. The channel contracted by about 121 acres in this reach since 1950, and about 150 acres of riparian vegetation has encroached into old channel areas. This pattern has been consistent in the lower river, and relates primarily to a reduction in flows due to human development. Floodplain turnover rates have dropped from 8.9 acres per year pre-1976 to 5.4 acres per year post-1976.

Even though no side channels have been intentionally blocked, Reach D7 has lost about 3,800 feet of side channel length since 1950. This is likely due to passive loss caused by a reduction in high flows. Lower flows have also resulted in the isolation of 48 percent of the historic 5-year floodplain.

Land use is predominantly agricultural, with about 258 acres of pivot irrigation development since 1950. There are 27 acres of pivot irrigation and 21 acres of exurban land uses in the Channel Migration Zone. Two dump sites have been mapped on the right bank at RM 84R and RM 85.9R.

There are 7.4 acres of mapped Russian olive in the reach.

Reach D7 was part of the avian study. A total of 43 species were identified in the reach, including the Ovenbird, which has been identified by the Montana Natural Heritage Program as a Potential Special Concern. The Black-billed Cuckoo and Red-headed Woodpecker were also identified, both of which are Species of Concern.

A hydrologic evaluation of flow depletions indicates that flow alterations over the last century have been major in this reach. The magnitude of the 100-year flood is now 127,000 cfs, which 12 percent lower than it was pre-development (145,000 cfs). The 2-year flood, which strongly influences overall channel form, has dropped by 22 percent. Low flows have also been impacted; severe low flows described as 7Q10 (the lowest average 7-day flow anticipated every ten years) for summer months has dropped from an estimated 4,700 cfs to 2,600 cfs with human development, a reduction of 45 percent. More typical summer low flows, described as the summer 95% flow duration, have dropped from 6,890 cfs under unregulated conditions to 3,110 cfs under regulated conditions, a reduction of 55 percent.

Seasonal low flows have increased by 78 percent in the winter and 62 percent in the fall. Both fall and winter base flows are currently about 3,500 cfs.

CEA-Related observations in Reach D7 include:


- Passive loss of side channels with flow alterations

Recommended Practices (may include Yellowstone River Recommended Practices--YRRPs) for Reach D7 include:

- Russian olive removal



## PHYSICAL FEATURES MAP (2011)



## HYDROLOGIC SUMMARY

Hydrologic data available for the Reach Narratives include data from representative gaging stations, modeling from the COE from the Big Horn river upstream, and modeling by the USGS for the Big Horn River to the Missouri River confluence. Gaging stations that best represent the watershed area within any reach are used to describe the flood history within the reach. Hydrology modeling results generated for all reaches provides unregulated and regulated flow values. Seasonal and annual flow duration data generated by the USGS are available for reaches C10 through D13.

### Gage Representation (Gage-Based): Sidney

#### Flood History

Year	Date	Flow on Date	Return Interval	Gage No	Downstream Gage	Upstream Gage
1978	May 23	111,000	10-25 yr	6329500	6309000	
1912	Mar 29	114,000	10-25 yr	Location	Sidney	Miles City
1944	Jun 21	120,000	10-25 yr	Period of Record	1911-2015	1929-2015
2011	May 24	124,000	10-25 yr	Distance To (miles)	50.6	95.0
1918	Jun 20	126,000	25-50 yr			
1943	Mar 29	132,000	25-50 yr			
1923	Oct 3	134,000	25-50 yr			
1952	Mar 31	138,000	25-50 yr			
1921	Jun 21	159,000	100-yr			

#### Discharge

	1.01 Yr	2 Yr	5 Yr	10 Yr	50 Yr	100 Yr	500 Yr	7Q10 Summer	95% Sum. Duration
Unregulated		69,500	89,600	103,000	133,000	145,000	176,000	4,700	6,890
Regulated		54,200	74,200	87,200	115,000	127,000	153,000	2,600	3,110
% Change		-22.01%	-17.19%	-15.34%	-13.53%	-12.41%	-13.07%	-44.68%	-54.86%

#### Flow Duration

Streamflow, in ft<sup>3</sup>/s, which was equaled or exceeded for indicated percent of time

Note that these statistics are only available from Reach C10 downstream. See the USGS report for detailed information.

Season		5%	50%	95%
Spring	Unregulated	67,500	25,100	6,960
	Regulated	52,100	14,900	5,080
	% Change	-23%	-41%	-27%
Summer	Unregulated	47,300	14,900	6,890
	Regulated	35,200	8,820	3,110
	% Change	-26%	-41%	-55%
Fall	Unregulated	9,800	5,940	2,010
	Regulated	11,200	7,430	3,570
	% Change	14%	25%	78%
Winter	Unregulated	14,800	5,380	2,120
	Regulated	15,400	6,550	3,440
	% Change	4%	22%	62%
Annual	Unregulated	49,900	8,900	2,820
	Regulated	37,200	8,020	3,620
	% Change	-25%	-10%	28%



## AERIAL PHOTOGRAPHY

A variety of aerial photographic sources provide the basis for much of the Cumulative Effects Assessment analysis. The table below lists the air photos compiled for the reach and the associated discharge at the most representative USGS gaging station.

	Source	Acquisition Date	Type	Scale	Gage	Discharge
1950	USGS-EROS	26-Aug-49	B/W	1:14,800	6329500	2750
1976	USCOE	9-Oct-76	B/W	1:24,000	6329500	9580
1995	USGS DOQQ	12-Jun-96	B/W		6329500	52600
2001	NRCS	August 2-8, 2001	CIR	1:24,000	6329500	4000
2004	Merrick	5/20/2004 - 6/3/04	Color	1:15,840	6329500	5070
2005	NAIP	07/14/2005	color	1-meter pixels	6329500	15900
2009	NAIP	8/10/2009	Color	1-meter pixels	6329500	13700
2009	NAIP	7/11/2009	Color	1-meter pixels	6329500	32600
2011	USCOE	October 2012	color	1-ft pixel	6329500	9030
2011	NAIP	7/20/2011	Color	1-meter pixels	6329500	48800
2013	NAIP	07/27/2013	color	1-meter pixels	6329500	

## PHYSICAL FEATURES

Several efforts to capture the types and extents of physical features in the corridor have been generated by the CEA study. The 2001 Physical Features Inventory was performed through helicopter/video Rapid Aerial Assessment by the NRCS (NRCS, 2001) and did not include Park County. This inventory includes point and linear features that represent bank armor, irrigation structures, transportation encroachments, and areas of accelerated erosion. Bank armor mapped in the 2001 inventory only reflects features on the active channel margin, and thus excludes off-channel features on historic side channels. Some floodplain restriction features such as dikes and levees in the 2001 Physical Features Inventory may extend well beyond the active channel. In 2013, the 2001 inventory was revised to include Park County. At that time, some attribute inconsistencies in the original data were addressed. This dataset was then updated to reflect conditions in the 2011 NAIP imagery.

For Stillwater, Yellowstone and Dawson Counties, a Physical Features Timeline was generated that includes additional mapping based on aerial photography and assigns approximate dates of feature construction based on observed presence/absence in historic imagery between the 1950s and 2005 (DTM and AGI, 2008). The Physical Features Timeline contains features that were not mapped in the 2001 inventory (e.g. bank armor abandoned in floodplain areas by 2001). As such the total bank armor extent in the 2005 data is commonly greater than that identified in 2001 or 2013.


**Note:** As the goal for each physical features mapping effort were different, with differing mapping extents, there will be discrepancies between total feature lengths (e.g. length of rock riprap) in each data set.

**Bankline/Floodplain Inventory: Time Series** The Human Impacts Timeline assessed physical feature development through time for Yellowstone, Stillwater, and Dawson Counties.

Feature Class	Feature Type	Sum of Feature Length (ft)					
		1950	1976	1995	2001	2004	2005
Transportation Encroachment	Railroad	12,529	12,529	12,529	12,529	12,529	12,529
	<b>Totals</b>	<b>12,529</b>	<b>12,529</b>	<b>12,529</b>	<b>12,529</b>	<b>12,529</b>	<b>12,529</b>

## ICE JAMS

Ice jam data were obtained from the National Ice Jam Database maintained by the Ice Engineering Group at Army Corps of Engineers Cold Regions Research and Engineering Laboratory (<https://rsgis.crrel.usace.army.mil/icejam/>). From this database, Yellowstone River ice jams are summarized by reach in the Yellowstone River Historic Events Timeline (DTM and AGI, 2008b). The basic information for each ice jam is presented as a list of events. The graph represents the number of database entries for a reach. Note that a single jam event may have multiple entries.



## GEOMORPHIC

The geomorphology data presented below consist of measured changes in Braiding Parameter since 1950 and blocked side channels. Braiding parameter is a measure of the total length of side channels relative to that of the main channel. The braiding parameter is calculated as the sum of anabranching and primary channel lengths divided by the primary channel length. Secondary channels within the bankfull margins are a function of flow stage and hence were not included in the braiding parameter calculation. If a reach has a braiding parameter of 3, then the total bankfull channel length is three times that of the main channel. The mean braiding parameter measured for all 88 reaches is 1.8.

Blocked side channels that were either plugged with a small dike or cutoff by larger features such as a levee or road prism were identified for the pre and post-1950s eras.

Additional geomorphic parameters are discussed in more detail in the study report and appendices.

### Braiding (Bankfull)

	Primary Chan. Length (ft)	Anab. Ch. Length (ft)	Bankfull Braiding Parameter		% Change in Braiding
1950	43,833	59,956	2.37	1950 to 1976:	1.78%
1976	39,713	55,991	2.41	1976 to 1995:	3.49%
1995	40,102	59,914	2.49	1995 to 2001:	-4.10%
2001	40,314	56,108	2.39	1950 to 2001:	1.01%
<b>Change 1950 - 2001</b>	<b>-3,519</b>	<b>-3,848</b>	<b>0.02</b>		

### Length of Side Channels Blocked

Pre-1950s (ft)	0
Post-1950s (ft)	0

## HYDRAULICS

Available hydraulic information includes county-based HEC-RAS modeling efforts by the Army Corps of Engineers with the exclusion of Park County. Floodplain modeling was performed for four conditions representing a developed and undeveloped floodplain, and unregulated and regulated flows for the 1.5, 2, 5, 10, 20, 50, 100, 200, and 500-year events. Park County has limited FEMA hydraulic modeling and was not included in the analysis.

The results of HEC-RAS modeling for the 5 and 100-year flood events were assessed to compare the extents of inundated area for the pristine (undeveloped floodplain, unregulated flows) and developed (developed floodplain, regulated flows) conditions. The data sets provided for each flow condition were unioned in the GIS to identify areas where the inundated extent differed. These areas of human-caused floodplain isolation due to either flow alterations or physical features such as levees. For the 100-year flood event, isolated areas greater than 5 acres were attributed with the interpreted reason for isolation (railroad, levee, etc.). The resulting values are presented as acres and percent of the pristine floodplain that has been isolated. The pristine floodplain is defined as the total floodplain footprint minus the area of the mapped 2001 bankfull channel (mapped islands were included in the floodplain area).

### Floodplain Isolation

	100-Year		5-Year	
	Isolated Acres	% of Floodplain	Isolated Acres	% of Floodplain
Non-Structural (hydrology, geomorphic, etc.)	44	2.4%		
Agriculture (generally relates to field boundaries)	0	0.0%		
Agriculture (isolated by canal or large ditch)	0	0.0%		
Levee/Riprap (protecting agricultural lands)	0	0.0%		
Levee/Riprap (protecting urban, industrial, etc.)	0	0.0%		
Railroad	0	0.0%		
Abandoned Railroad	0	0.0%		
Transportation (Interstate and other roads)	0	0.0%		
<b>Total Not Isolated (Ac)</b>	1762		1532	
<b>Total Floodplain Area (Ac)</b>	1806		1928	
<b>Total Isolated (Ac)</b>	<b>44</b>	<b>2.4%</b>	<b>395</b>	<b>47.9%</b>

The 5-year floodplain is a good allegory for the extent of the riparian zone. Thus, irrigated areas within the 5-year floodplain tend to represent riparian zones that have been converted to agriculture and may result in additional bank protection to protect the agricultural production and irrigation infrastructure.

	Flood	Sprinkler	Pivot	Total
Irrigated Acres within the 5 Year Flooplain:	2	0	0	2

## CHANNEL MIGRATION ZONE

A series of Channel Migration Maps were developed for the Yellowstone River from Gardiner to its mouth in McKenzie County, North Dakota (Thatcher, Swindell, and Boyd, 2009). These maps and their accompanying report can be accessed from the YRCDC Website. The channel migration zone (CMZ) developed for the Yellowstone River is defined as a composite area made up of the existing channel, the historic channel since 1950 (Historic Migration Zone, or HMZ), and an Erosion Buffer that encompasses areas prone to channel erosion over the next 100 years. Areas within this CMZ that have been isolated by constructed features such as armor or floodplain dikes are attributed as “Restricted Migration Areas” (RMA). Beyond the CMZ boundaries, outlying areas that pose risks of channel avulsion are identified as “Avulsion Potential Zones”.

Mean 50-Yr Migration Distance (ft)	Erosion Buffer (ft)	Total CMZ Acreage	Restricted CMZ Acreage	% Restricted Migration Area	Total AHZ Acreage	Restricted AHZ Acreage	% Restricted Avulsion Area
341	682	2,811	6	0%	127	0	0%

### 2011 Restricted Migration Area Summary

Note that these data reflect the observed conditions in the 2011 aerial photography (NAIP for Park and Sweet Grass Counties, COE for the rest of the river).

Reason for Restriction	Land Use Protected	RMA Acres	Percent of CMZ
Road/Railroad Prism			
	Railroad	6	0.2%
<b>Totals</b>		<b>6</b>	<b>0.2%</b>

### Land Uses within the CMZ (Acres)

Flood Irrigation	Sprinkler Irrigation	Pivot Irrigation	Urban/ExUrban	Transportation
180.4	2.2	27.3	20.7	9.0



## LAND USE

Land uses were mapped from aerial photography Gardiner to the confluence of the Missouri River in North Dakota for four time periods: 1950s, 1976, 2001, and 2011. Mapping was performed at approximately 1:6,000 to ensure consistent mapping across all data sets. Typically, if a feature could not be easily mapped at the target mapping scale, it was not separated out from the adjacent land use.

A four-tiered system was used to allow analysis at a variety of levels. Tier 1 breaks land use into Agricultural and Non-Agricultural uses. Tier two subdivided uses into productive Agricultural Land and Infrastructure for the Agricultural land, and Urban, Exurban and Transportation categories for the Non-Agricultural land. Tier three further breaks down land uses into more refined categories such as Irrigated or Non-Irrigated and Residential, Commercial, or Industrial. Finally, Tier 4 focuses primarily on the productive agricultural lands, identifying the type of irrigation (Pivot, Sprinkler or Flood).

### Land Use Timeline - Tiers 2 and 3

Feature Class	Feature Type	Acres				% of Reach Area			
		1950	1976	2001	2011	1950	1976	2001	2011
Agricultural Infrastructure									
	Canal	0	0	0	0	0.0%	0.0%	0.0%	0.0%
	Agricultural Roads	0	0	0	0	0.0%	0.0%	0.0%	0.0%
	Other Infrastructure	29	64	81	84	0.4%	0.9%	1.2%	1.2%
	<b>Totals</b>	<b>29</b>	<b>64</b>	<b>81</b>	<b>84</b>	<b>0.4%</b>	<b>0.9%</b>	<b>1.2%</b>	<b>1.2%</b>
Agricultural Land									
	Non-Irrigated	4,756	4,485	3,767	3,629	70.5%	66.5%	55.9%	53.8%
	Irrigated	0	182	876	992	0.0%	2.7%	13.0%	14.7%
	<b>Totals</b>	<b>4,756</b>	<b>4,668</b>	<b>4,644</b>	<b>4,621</b>	<b>70.5%</b>	<b>69.2%</b>	<b>68.9%</b>	<b>68.5%</b>
Channel									
	Channel	1,869	1,918	1,881	1,899	27.7%	28.4%	27.9%	28.2%
	<b>Totals</b>	<b>1,869</b>	<b>1,918</b>	<b>1,881</b>	<b>1,899</b>	<b>27.7%</b>	<b>28.4%</b>	<b>27.9%</b>	<b>28.2%</b>
ExUrban									
	ExUrban Other	0	0	24	23	0.0%	0.0%	0.4%	0.3%
	ExUrban Undeveloped	0	0	0	0	0.0%	0.0%	0.0%	0.0%
	ExUrban Industrial	0	0	0	0	0.0%	0.0%	0.0%	0.0%
	ExUrban Commercial	0	0	0	0	0.0%	0.0%	0.0%	0.0%
	ExUrban Residential	0	3	22	26	0.0%	0.0%	0.3%	0.4%
	<b>Totals</b>	<b>0</b>	<b>3</b>	<b>46</b>	<b>49</b>	<b>0.0%</b>	<b>0.0%</b>	<b>0.7%</b>	<b>0.7%</b>
Transportation									
	Public Road	57	59	59	59	0.8%	0.9%	0.9%	0.9%
	Interstate	0	0	0	0	0.0%	0.0%	0.0%	0.0%
	Railroad	31	31	31	31	0.5%	0.5%	0.5%	0.5%
	<b>Totals</b>	<b>88</b>	<b>90</b>	<b>90</b>	<b>90</b>	<b>1.3%</b>	<b>1.3%</b>	<b>1.3%</b>	<b>1.3%</b>
Urban									
	Urban Other	0	0	0	0	0.0%	0.0%	0.0%	0.0%
	Urban Residential	0	0	0	0	0.0%	0.0%	0.0%	0.0%
	Urban Commercial	0	0	0	0	0.0%	0.0%	0.0%	0.0%
	Urban Undeveloped	0	0	0	0	0.0%	0.0%	0.0%	0.0%
	Urban Industrial	0	0	0	0	0.0%	0.0%	0.0%	0.0%
	<b>Totals</b>	<b>0</b>	<b>0</b>	<b>0</b>	<b>0</b>	<b>0.0%</b>	<b>0.0%</b>	<b>0.0%</b>	<b>0.0%</b>

### Land Use Timeline - Tiers 3 and 4

Feature Class	Feature Type	Acres				% of Reach Area				Change Between Years (% of Agricultural Land)			
		1950	1976	2001	2011	1950	1976	2001	2011	'50-76	'76-01	'01-11	'50-11
Irrigated													
	Sprinkler	0	0	26	26	0.0%	0.0%	0.5%	0.6%	0.0%	0.5%	0.0%	0.6%
	Pivot	0	0	0	258	0.0%	0.0%	0.0%	5.6%	0.0%	0.0%	5.6%	5.6%
	Flood	0	182	851	708	0.0%	3.9%	18.3%	15.3%	3.9%	14.4%	-3.0%	15.3%
	<b>Totals</b>	<b>0</b>	<b>182</b>	<b>876</b>	<b>992</b>	<b>0.0%</b>	<b>3.9%</b>	<b>18.9%</b>	<b>21.5%</b>	<b>3.9%</b>	<b>15.0%</b>	<b>2.6%</b>	<b>21.5%</b>

Non-Irrigated

Multi-Use	3,714	2,925	2,899	2,816	78.1%	62.7%	62.4%	61.0%	-15.4%	-0.2%	-1.5%	-17.1%
Hay/Pasture	1,043	1,560	868	812	21.9%	33.4%	18.7%	17.6%	11.5%	-14.7%	-1.1%	-4.3%
<b>Totals</b>	<b>4,756</b>	<b>4,485</b>	<b>3,767</b>	<b>3,629</b>	<b>#####</b>	<b>96.1%</b>	<b>81.1%</b>	<b>78.5%</b>	<b>-3.9%</b>	<b>-15.0%</b>	<b>-2.6%</b>	<b>-21.5%</b>

## RIPARIAN

Riparian mapping data are derived from the Yellowstone River Riparian Vegetation Mapping study (DTM/AGI 2008). This study coarsely mapped the riparian vegetation communities using 1950's, 1976-1977, and 2001 aerial imagery in a GIS environment. The polygons are digitized at a scale of approximately 1:7,500, with a minimum mapping unit of approximately 10 acres. The goal of the delineation was to capture areas of similar vegetation structure as they appeared on the aerial imagery, while maintaining a consistent scale.

The "Riparian Turnover" values quantify the total area within the active channel area that converted from either woody vegetation to open bar or water, or from open bar or water to woody vegetation. A comparison of these values allows some consideration of overall riparian encroachment into the river corridor from 1950 to 2001.

### Riparian Mapping

Statistic	Shrub (Acres)			Closed Timber (Acres)			Open Timber (Acres)		
	1950	1976	2001	1950	1976	2001	1950	1976	2001
Min	0.2	0.5	0.9	0.1	0.7	2.7	2.7	2.0	1.4
Max	107.9	88.8	35.3	153.3	159.7	298.9	53.7	26.1	48.2
Average	13.4	13.5	8.2	44.6	35.5	66.1	15.1	11.3	16.8
Sum	617.6	619.7	318.1	757.7	815.5	1,123.3	136.3	67.6	134.1

### Riparian Turnover

Conversion of riparian areas to channel, or from channel to riparian between the 1950's and 2001 data set.

Riparian to Channel (acres) 209.9

Channel to Riparian (acres) 359.2

**Riparian Encroachment (acres) 149.4**

### Riparian Recruitment

Creation of riparian areas between 1950s and 2001.	1950s Channel Mapped as 2011 Riparian (Ac)	366.8
	1950s Floodplain Mapped as 2011 Channel (Ac)	53.7
	<b>Total Recruitment (1950s to 2011)(Ac)</b>	<b>420.6</b>

## WETLANDS

Wetland areas were mapped to National Wetland Inventory standards by the Montana Natural Heritage Program. Palustrine wetlands within the mapped 100-year inundation boundary were extracted and summarized into four categories: Riverine (Unconsolidated Bottom - UB, Aquatic Bed - AB, and Unconsolidated Shore - US), Emergent - EM, Scrub-Shrub - SS, and Forested - FO.

	Riverine	Emergent	Scrub/Shrub	Forested	Total
<b>Mapped Acres</b>	28.9	72.3	47.1	0.0	<b>148.2</b>
<b>Acres/Valley Mile</b>	4.2	10.6	6.9	0.0	

## RUSSIAN OLIVE

Russian olive is considered an invasive species and its presence in the Yellowstone River corridor is fairly recent. As such, its spread can be used as a general indicator of invasive plants within the corridor. It has the added benefit of being easily identified in multi-spectral aerial photography, making it possible to inventory large areas using remote techniques.

In 2011, Natural Resources Conservation Service (NRCS) in Bozeman, MT conducted an inventory of Russian olive locations in the Yellowstone River watershed. This study utilized the Feature Analyst extension within ArcGIS to interpret multi-spectral 2008 NAIP imagery for the presence of Russian olive. The resulting analysis was converted from raster format to a polygon ESRI shape file for distribution and further analysis within a GIS environment.

This work scope was tasked with integrating the resulting Russian olive inventory into the Yellowstone River Conservation Districts Council (YRDC) Cumulative Effects Assessment (CEA) GIS and associated reach-based database. Additionally, analysis of Russian olive within the corridor was conducted to characterize its distribution in throughout the corridor and its association with other corridor data sets.

	Floodplain Area (Ac)	% of Floodplain	Other Area (Ac)	Inside RMA (Ac)	Inside '50s Channel (Ac)	Inside 50s Island (Ac)
<b>Russian Olive in Reach</b>	7.44	0.21%	1.67	0.00	4.97	1.12

## FISHERIES SUMMARY

Fisheries data available for the Reach Narratives include low-flow and high-flow habitat mapping of 2001 conditions for 406 miles of river, extending from the mouth upstream to a point approximately 8 miles upstream of Park City. Habitat mapping was performed remotely on the 2001 CIR aerial photography utilizing habitat classifications developed by Montana Fish, Wildlife, and Parks (DTM 2009). Historic habitat mapping using the 1950's imagery is limited to Reach B1 (high-flow) and D9 (low and high-flow).

Fisheries field sampling data have been provided by Ann Marie Reinhold (MSU). In this study, the Yellowstone River from Park City to Sidney was divided into five segments. Within each segment, fish were sampled in reaches modified by riprap ("treatment reaches") and relatively unmodified reaches ("control reaches"). Fish sampling was conducted during summer and autumn of 2009, 2010, and 2011. Boat electrofishing, trammel nets, mini-fyke nets and bag seines were used to collect data from river bends.

Fish presence data is only presented for those reaches that were sampled.

The Low Flow Habitat Mapping followed schema developed by Montana Fish Wildlife and Parks to identify key habitat units for certain aquatic species.

### Low Flow Fisheries Habitat Mapping

Habitat	2001 (Acres)		
	Bankfull	Low Flow	% of Low Flow
Scour Pool	349.7	212.3	11.3%
Bluff Pool	138.1	163.2	8.7%
Secondary Channel	186.2	130.8	7.0%
Secondary Channel (Seasonal)	262.8	190.1	10.1%
Channel Crossover	164.3	124.7	6.6%
Point Bar		89.5	4.8%
Side Bar		69.3	3.7%
Mid-channel Bar		60.2	3.2%
Island	778.5	789.4	42.0%
Dry Channel		47.6	2.5%

## AVIAN

Birds were sampled in 2006 and 2007 by Danielle Jones of Montana State University. Point count methods were used at 304 randomly chosen sites in 21 braided or anabranching reaches. Each site was visited multiple times within a season, and sites were visited in both years. Birds were sampled in grassland, shrubland, and cottonwood forest habitats. Additional bird data was collected by Amy Cilimburg of Montana Audubon in summer 2012. High priority areas for data collection were identified with the assistance of the YRCDC Technical Advisory Committee. The Audubon methodology recorded data for a wider variety of bird species relative to the MSU study, including raptors and waterfowl.

Bird Species Observed in Reach/Region		Species of Concern	Potential Species of Concern
<input checked="" type="checkbox"/> <input checked="" type="checkbox"/> American Robin	<input checked="" type="checkbox"/> <input checked="" type="checkbox"/> Chipping Sparrow	<input checked="" type="checkbox"/> <input checked="" type="checkbox"/> Killdeer	<input checked="" type="checkbox"/> <input checked="" type="checkbox"/> Song Sparrow
<input checked="" type="checkbox"/> <input checked="" type="checkbox"/> American Crow	<input type="checkbox"/> <input checked="" type="checkbox"/> Clay-collared Sparrow	<input type="checkbox"/> <input type="checkbox"/> Lark Bunting	<input checked="" type="checkbox"/> <input checked="" type="checkbox"/> Spotted Sandpiper
<input checked="" type="checkbox"/> <input checked="" type="checkbox"/> American Goldfinch	<input type="checkbox"/> <input type="checkbox"/> Cliff Swallow	<input checked="" type="checkbox"/> <input checked="" type="checkbox"/> Lark Sparrow	<input checked="" type="checkbox"/> <input checked="" type="checkbox"/> Spotted Towhee
<input checked="" type="checkbox"/> <input checked="" type="checkbox"/> American Kestrel	<input checked="" type="checkbox"/> <input checked="" type="checkbox"/> Common Grackle	<input checked="" type="checkbox"/> <input checked="" type="checkbox"/> Lazuli Bunting	<input type="checkbox"/> <input type="checkbox"/> Sharp-shinned Hawk
<input checked="" type="checkbox"/> <input checked="" type="checkbox"/> American Redstart	<input type="checkbox"/> <input type="checkbox"/> Common Merganser	<input checked="" type="checkbox"/> <input checked="" type="checkbox"/> Least Flycatcher	<input type="checkbox"/> <input type="checkbox"/> Swainson's Thrush
<input checked="" type="checkbox"/> <input checked="" type="checkbox"/> Bald Eagle	<input type="checkbox"/> <input checked="" type="checkbox"/> Common Nighthawk	<input type="checkbox"/> <input checked="" type="checkbox"/> Mallard	<input type="checkbox"/> <input type="checkbox"/> Sandhill Crane
<input type="checkbox"/> <input type="checkbox"/> Baltimore Oriole	<input type="checkbox"/> <input type="checkbox"/> Common Raven	<input type="checkbox"/> <input type="checkbox"/> Mountain Bluebird	<input checked="" type="checkbox"/> <input checked="" type="checkbox"/> Tree Swallow
<input type="checkbox"/> <input type="checkbox"/> Barn Swallow	<input checked="" type="checkbox"/> <input checked="" type="checkbox"/> Common Yellowthroat	<input checked="" type="checkbox"/> <input checked="" type="checkbox"/> Mourning Dove	<input type="checkbox"/> <input type="checkbox"/> Turkey Vulture
<input type="checkbox"/> <input checked="" type="checkbox"/> Belted Kingfisher	<input type="checkbox"/> <input checked="" type="checkbox"/> Cooper's Hawk	<input checked="" type="checkbox"/> <input checked="" type="checkbox"/> Northern Flicker	<input type="checkbox"/> <input type="checkbox"/> Upland Sandpiper
<input checked="" type="checkbox"/> <input checked="" type="checkbox"/> Black-billed Cuckoo	<input type="checkbox"/> <input checked="" type="checkbox"/> Dickcissel	<input checked="" type="checkbox"/> <input checked="" type="checkbox"/> Orchard Oriole	<input type="checkbox"/> <input type="checkbox"/> Vesper Sparrow
<input checked="" type="checkbox"/> <input checked="" type="checkbox"/> Black-billed Magpie	<input checked="" type="checkbox"/> <input checked="" type="checkbox"/> Downy Woodpecker	<input type="checkbox"/> <input type="checkbox"/> Osprey	<input type="checkbox"/> <input type="checkbox"/> Violet-green Swallow
<input checked="" type="checkbox"/> <input checked="" type="checkbox"/> Black-capped Chickadee	<input type="checkbox"/> <input checked="" type="checkbox"/> Eastern Bluebird	<input checked="" type="checkbox"/> <input checked="" type="checkbox"/> Ovenbird	<input checked="" type="checkbox"/> <input checked="" type="checkbox"/> Warbling Vireo
<input type="checkbox"/> <input checked="" type="checkbox"/> Black-and-white Warbler	<input checked="" type="checkbox"/> <input checked="" type="checkbox"/> Eastern Kingbird	<input type="checkbox"/> <input checked="" type="checkbox"/> Plumbeous Vireo	<input checked="" type="checkbox"/> <input checked="" type="checkbox"/> Western Kingbird
<input checked="" type="checkbox"/> <input checked="" type="checkbox"/> Black-headed Grosbeak	<input type="checkbox"/> <input type="checkbox"/> Eurasian Collared-dove	<input checked="" type="checkbox"/> <input checked="" type="checkbox"/> Red-headed Woodpecker	<input checked="" type="checkbox"/> <input checked="" type="checkbox"/> Western Meadowlark
<input type="checkbox"/> <input checked="" type="checkbox"/> Blue Jay	<input checked="" type="checkbox"/> <input checked="" type="checkbox"/> European Starling	<input type="checkbox"/> <input type="checkbox"/> Red-naped Sapsucker	<input checked="" type="checkbox"/> <input checked="" type="checkbox"/> Western Wood-pewee
<input type="checkbox"/> <input checked="" type="checkbox"/> Bobolink	<input type="checkbox"/> <input checked="" type="checkbox"/> Field Sparrow	<input type="checkbox"/> <input checked="" type="checkbox"/> Red Crossbill	<input checked="" type="checkbox"/> <input checked="" type="checkbox"/> White-breasted Nuthatch
<input checked="" type="checkbox"/> <input checked="" type="checkbox"/> Brewer's Blackbird	<input type="checkbox"/> <input checked="" type="checkbox"/> Franklin's Gull	<input checked="" type="checkbox"/> <input checked="" type="checkbox"/> Ring-necked Pheasant	<input type="checkbox"/> <input type="checkbox"/> White-throated Swift
<input checked="" type="checkbox"/> <input checked="" type="checkbox"/> Brown-headed Cowbird	<input type="checkbox"/> <input checked="" type="checkbox"/> Grasshopper Sparrow	<input type="checkbox"/> <input checked="" type="checkbox"/> Red-tailed hawk	<input type="checkbox"/> <input checked="" type="checkbox"/> Wild Turkey
<input type="checkbox"/> <input type="checkbox"/> Brown Creeper	<input type="checkbox"/> <input checked="" type="checkbox"/> Gray Catbird	<input type="checkbox"/> <input checked="" type="checkbox"/> Rock Dove	<input type="checkbox"/> <input checked="" type="checkbox"/> Wood Duck
<input type="checkbox"/> <input checked="" type="checkbox"/> Brown Thrasher	<input type="checkbox"/> <input checked="" type="checkbox"/> Great Blue Heron	<input type="checkbox"/> <input checked="" type="checkbox"/> Red-winged Blackbird	<input type="checkbox"/> <input checked="" type="checkbox"/> Yellow-bellied Sapsucker
<input type="checkbox"/> <input checked="" type="checkbox"/> Bullock's Oriole	<input type="checkbox"/> <input checked="" type="checkbox"/> Great Horned Owl	<input checked="" type="checkbox"/> <input checked="" type="checkbox"/> Red-eyed Vireo	<input type="checkbox"/> <input checked="" type="checkbox"/> Yellow-billed Cuckoo
<input type="checkbox"/> <input type="checkbox"/> Canada Goose	<input checked="" type="checkbox"/> <input checked="" type="checkbox"/> Hairy Woodpecker	<input type="checkbox"/> <input checked="" type="checkbox"/> Red-breasted Grosbeak	<input checked="" type="checkbox"/> <input checked="" type="checkbox"/> Yellow-breasted Chat
<input checked="" type="checkbox"/> <input checked="" type="checkbox"/> Cedar Waxwing	<input type="checkbox"/> <input type="checkbox"/> House Finch	<input type="checkbox"/> <input type="checkbox"/> Say's Phoebe	<input type="checkbox"/> <input type="checkbox"/> Yellow-headed Blackbird
<input type="checkbox"/> <input checked="" type="checkbox"/> Chimney Swift	<input checked="" type="checkbox"/> <input checked="" type="checkbox"/> House Wren	<input type="checkbox"/> <input checked="" type="checkbox"/> Savannah Sparrow	<input checked="" type="checkbox"/> <input checked="" type="checkbox"/> Yellow Warbler



## CULTURAL INVENTORY SUMMARY

The Yellowstone River Cultural Inventory - 2006 documents the variety and intensity of different perspectives and values held by people who share the Yellowstone River. Between May and November of 2006, a total of 313 individuals participated in the study. They represented agricultural, civic, recreational, or residential interest groups. Also, individuals from the Crow and the Northern Cheyenne tribes were included.

There are three particular goals associated with the investigation. The first goal is to document how the people of the Yellowstone River describe the physical character of the river and how they think the physical processes, such as floods and erosion, should be managed. Within this goal, efforts have been made to document participants' views regarding the many different bank stabilization techniques employed by landowners. The second goal is to document the degree to which the riparian zone associated with the river is recognized and valued by the participants. The third goal is to document concerns regarding the management of the river's resources. Special attention is given to the ways in which residents from diverse geographical settings and diverse interest groups view river management and uses. The results illustrate the commonalities of thought and the complexities of concerns expressed by those who share the resources of the Yellowstone River.

### Summary of Cultural Views in Region D

A review of the interview data for the segment, Missouri River to Powder River, suggests that people in this area engage in four primary discussions when asked about the Yellowstone River. First, the notion of Eastern Montana is not simply a geographic reference. It is a defining concept that captures the agricultural roots and the cultural values of the people living in the study segment, and the river is an essential element within their notion of Eastern Montana. Second, the river is discussed as a wholesome recreational outlet. However, shifting landownership is noted as an important change in the recreational context. Third, even though agricultural practices are viewed as the mainstay of the local economies, many participants discuss the long-term economic viability of their communities as a concern. Industrial and residential developments along the river's edge are seemingly remote possibilities and are generally discussed with references to flood plain restrictions and the stability of nearby dikes. Finally, discussions of managing the river are limited, but a variety of opinions are offered regarding bank erosion and stabilization techniques.