

| | | | |
|------------------|--------------------------------------|-----------------------|--------------------|
| County | Stillwater | Upstream River Mile | 413.7 |
| Classification | PCA: Partially confined anabranching | Downstream River Mile | 405.9 |
| General Location | Below Columbus | Length | 7.80 mi (12.55 km) |

Narrative Summary

Reach A14 is located in Stillwater County, just downstream of Columbus. The reach is a Partially Confined Anabranching (PCA) reach type, reflecting some valley while influence coupled with relatively extensive forested islands. The reach is 7.8 miles long, extending from RM 405.9 to RM 413.7. The partial geologic confinement within Reach A14 is created by interbedded sandstone and shale of the Cretaceous-age Judith River Formation that intermittently forms the active channel margin on either its right or left bank. The Parkman Sandstone, a massive cliff-forming unit within the Judith River Formation, forms cliffs against the channel that are commonly over 150 feet high.

Similar to other reaches in Region A, the overall footprint of the river channel has increased in size since 1950. In 1950, the channel footprint was 637 acres but by 2001 it had expanded to 728 acres. This was accompanied by a net loss of about 32 acres of riparian area to channel during that same timeframe.

Approximately 16 percent of the bankline in Reach A14 is armored, and the armor is almost entirely rock riprap, with a very short section of flow deflectors. The armor is located almost entirely on the northern corridor margin, where transportation infrastructure (mainly railroad) follows the edge of the valley.

Over three miles of side channels have been blocked in Reach A14, with about half of the blockages occurring prior to 1950 and half after. The losses occurred on two distinct channels, one at RM 410 on the south side of the corridor and one at RM 407 on the north side.

Land use in Reach A14 is almost entirely agricultural, with almost 260 acres mapped as agricultural infrastructure. This in part reflects corrals that are part of an animal handling facility on the north side of the river at RM 409. There are 1,300 acres under flood irrigation in the reach, and 144 acres in pivot. A total of 227 acres of developed land are in the Channel Migration Zone, most of that is in flood irrigation (215 acres). Less than 2 percent of the CMZ is isolated by physical features, all of which is behind the armored rail line on the north side of the river.

There is one major diversion in Reach A14; Cove Ditch diverts water from the north bank at RM 410.

Reach A14 was sampled as part of the avian study. The average species richness in Reach A14 was 7.9, which indicates the average number of species observed during site visits to the reach in cottonwood habitats. The average species richness for all sites evaluated is 8. Riparian mapping in Reach A14 shows a reduction of about 100 acres of closed timber in the reach since 1950. Since 1950, Reach A14 has lost most of its forest that would be considered at low risk of cowbird infestation due to its separation from agricultural infrastructure. In 1950, about 10.5 acres of forest per valley mile were identified as low risk and by 2001 that forest area had been reduced to 0.5 acres per valley mile.

Reach A14 has approximately 2.5 acres of mapped Russian olive, which is concentrated along ditches and low riparian/wetland areas north of the river. There are also over 250 acres of mapped wetland in the reach, most of which is emergent marshes and wet meadows. About 27 acres of emergent wetland have been isolated from the river corridor by the rail line at RM 413.5.

A hydrologic evaluation of flow depletions indicates that flow alterations over the last century have been moderate in this reach. The mean annual flood is estimated to have dropped from 16,200 cfs to 15,100 cfs, a drop of about 7 percent. The biggest influence has been on low flows: severe low flows described as 7Q10 (the lowest average 7-day flow anticipated every ten years) for summer months has dropped from an estimated 2,280 cfs to 1,770 cfs with human development, a reduction of 22 percent. More typical summer low flows, described as the summer 95% flow duration, have dropped from 1,760 cfs under unregulated conditions to 1,680 cfs under regulated conditions at the Livingston gage, a reduction of 4.6 percent.

CEA-Related observations in Reach A14 include:

- Isolation of large wetland area by rail line
- Over 3 miles of side channel blockages
- Large corrals that are part of an animal handling facility within 1,000 feet of the riverbank


Recommended Practices (may include Yellowstone River Recommended Practices--YRRPs) for Reach A14 include:

- Side channel restoration at RM 410 and RM 407
- Russian olive removal (2.5 acres)
- Nutrient management at corrals that are part of an animal handling facility at RM 409
- Irrigation diversion structure management at Cove Ditch Diversion
- Wetland management/restoration at large complex isolated from river by rail line at RM 413.5


The following table summarizes some key CEA results that have been used to describe overall condition and types of human influences affecting the river. The values are specific to this single reach. Blanks indicate that a particular value was not available for this area. This information is consolidated from a large dataset that is presented in more detail in the full reach narrative report.

| | | | | | | |
|---|-------------------------|----------------------------|---|--|---|---|
| Discharge | Undev. | Developed | % Change | "Undeveloped" flows represent conditions prior to significant human development, whereas "developed" flows reflect the current condition of both consumptive and non-consumptive water use. | | |
| 2 Year (cfs) | 31,000 | 29,800 | -3.9% | | | |
| 100 Year (cfs) | 56,600 | 55,900 | -1.2% | | | |
| Bankfull Channel Area (Ac) | 1950 | 1976 | 1995 | 2001 | 1950-2001 | Bankfull channel area is the total footprint of the river inundated at approx. the 2-year flood. |
| | 637.3 | 675.2 | 635.5 | 727.9 | 90.6 | |
| Physical Features | 2011 Length (ft) | % of Bankline | 2001-2011 Change | There are additional types of bank armor such as car bodies and steel retaining walls, but they are relatively minor. | | |
| Rock RipRap | 13,457 | 16.4% | 1,807 | | | |
| Concrete Riprap | 0 | 0.0% | 0 | | | |
| Flow Deflectors | 64 | 0.1% | 0 | | | |
| Total | 13,521 | 16.5% | 1,807 | | | |
| Length of Side Channels Blocked (ft) | Pre-1950s | Post-1950s | Numerous side channels have been blocked by small dikes. | | | |
| | 9,672 | 9,176 | | | | |
| Floodplain Turnover | 1950 - 1976 | 1976 - 2001 | 1950-2001 In-channel riparian encroachment (negative number indicates retreat) | The rate of floodplain turnover reflects how many acres of land are eroded by the river. Turnover is associated with the creation of riparian habitat. | | |
| Total Acres | 185.7 | 141.7 | -31.84 acres | | | |
| Acres/Year | 7.1 | 5.7 | | | | |
| Acres/Year/Valley Mile | 1.0 | 0.8 | | | | |
| Open Bar Area | Point Bars | Bank Attached | Mid-Channel | Total | The type and extent of open sand and gravel bars reflect in-stream habitat conditions that can be important to fish, amphibians, and ground-nesting birds such as least terns. | |
| Change in Area '50 - '01 (Ac) | | | | | | |
| Floodplain Isolation | Acres | % of FP | Floodplain isolation refers to area that historically was flooded, but has become isolated do to flow alterations or physical features such as levees. | | | |
| 5 Year | 40.7 | 13% | | | | |
| 100 Year | 0.0 | 0% | | | | |
| Restricted Migration Area | Acres | % of CMZ | Channel Migration Zone restrictions refer to the area and percent of the CMZ that has been isolated by features such as bank armor, dikes, levees, and transportation embankments. | | | |
| | 25.7 | 1% | | | | |
| Land Use | 1950 | 2011 | 1950 | 2011 | Changes in land use reflect the development of the river corridor through time. The irrigated agricultural are is a sub-set of the mapped agricultural land. | |
| Agricultural Land (Ac) | 4,716.0 | 4,443.6 | Flood (Ac) | 1,663.6 | 1,319.8 | |
| Ag. Infrastructure (Ac) | 73.7 | 258.5 | Sprinkler (Ac) | 0.0 | 0.0 | |
| Exurban (Ac) | 0.0 | 0.0 | Pivot (Ac) | 0.0 | 144.0 | |
| Urban (Ac) | 0.0 | 0.0 | | | | |
| Transportation (Ac) | 90.2 | 188.5 | | | | |
| 1950s Riparian Vegetation Converted to a Developed Land Use (ac) | To Irrigated | To Other Use | Total Rip. Converted | % of 1950s Rip. | Changes in the extents of riparian vegetation are influenced by land use changes within the corridor. | |
| | 11.7 | 3.2 | 14.9 | 2.0% | | |
| National Wetlands Inventory | Acres | Acres per Valley Mi | Total Wetland Acres | Wetlands units summarized from National Wetlands Inventory Mapping include Riverine (typically open water sloughs), Emergent (marshes and wet meadows) and Shrub-Scrub (open bar areas with colonizing woody vegetation). | | |
| Riverine | 14.4 | 2.0 | 283.3 | | | |
| Emergent | 211.3 | 29.3 | | | | |
| Scrub/Shrub | 57.6 | 8.0 | | | | |
| Russian Olive (2001) (Appx. 100-yr Floodplain) | Acres | % | Russian olive is considered an invasive species and its presence in the corridor is fairly recent. Its spread can be used as a general indicator of invasive plants within the corridor. | | | |
| | 2.5 | 0.1% | | | | |
| Riparian Forest at low risk of Cowbird Parasitism (Ac/Valley Mile) | 1950 | 1976 | 2001 | Change 1950-2011 | Cowbirds are associated with agricultural and residential development, displacing native bird species by parasitizing their nests. | |
| | 10.5 | 0.5 | 0.5 | -10.0 | | |

PHYSICAL FEATURES MAP (2011)



CHANNEL MIGRATION ZONE MAP



| | | | |
|------------------|---------------------------------------|-----------------------|-------------------|
| County | Stillwater | Upstream River Mile | 405.9 |
| Classification | PCB: Partially confined braided | Downstream River Mile | 400 |
| General Location | Follows Stillwater/Carbon County line | Length | 5.90 mi (9.50 km) |

Narrative Summary

Reach A15 is located in Stillwater County between Columbus and Park City. The reach is a Partially Confined Braided (PCB) reach type, reflecting some valley wall influence coupled with relatively extensive open gravel bars and low flow channels. The reach is 5.9 miles long. The partial geologic confinement within Reach A15 is created by interbedded sandstone and shale of the Cretaceous-age Judith River Formation that intermittently forms the active channel margin on its right bank. The Parkman Sandstone, a massive cliff-forming unit within the Judith River Formation, forms cliffs against the channel that are commonly over 150 feet high.

Approximately 8 percent of the bankline in Reach A15 is armored, and the armor is almost entirely rock riprap, with a very short section of concrete armor. The armor is entirely located on the north bank of the river, across from the bluffs to the south.

Although no side channels have been mapped as blocked in the reach, the total anabranching channel length has dropped from 6.2 miles in 1950 to 4.2 miles in 2001.

Land use in Reach A15 is almost entirely agricultural, with over 200 acres mapped as agricultural infrastructure. This includes a large corral complex that is part of an animal handling facility on the north side of the river at RM 404. The corrals are behind a canal, but within a few hundred feet of the riverbank. There are 528 acres under flood irrigation in the reach, and 81 acres in pivot. A total of 119 acres of developed land are in the Channel Migration Zone, and all of that land is in flood irrigation. About 9 percent of the CMZ is isolated by physical features, all of which is behind armored canals associated with the Big Ditch Diversion, which diverts water from the north bank at RM 405.3. The Big Ditch Diversion structure fully spans a side channel of the river that is about 275 feet wide.

Riparian mapping in Reach A15 shows a reduction of about 60 acres of closed timber in the reach since 1950. Riparian recruitment rates have been relatively high; between 1950 and 2001 there were 200 acres of areas that recruited new riparian vegetation, and most of that was in old 1950s channels that were abandoned and became colonized. These abandoned channels also have high concentrations of Russian olive. Since 1950, Reach A15 has lost almost all of its forest that would be considered at low risk of cowbird infestation due to its separation from agricultural infrastructure. In 1950, about 20 acres of forest per valley mile were identified as low risk and by 2001 that forest area had been reduced to 1.

There are also over 150 acres of mapped wetland in the reach, most of which is emergent marshes and wet meadows. Large expanses of emergent wetlands have developed in side channels that have been passively lost since 1950 ("passively" meaning not blocked but abandoned).

A hydrologic evaluation of flow depletions indicates that flow alterations over the last century have been moderate in this reach. The mean annual flood is estimated to have dropped from 16,200 cfs to 15,100 cfs, a drop of about 7 percent. The biggest influence has been on low flows: severe low flows described as 7Q10 (the lowest average 7-day flow anticipated every ten years) for summer months has dropped from an estimated 2,286 cfs to 1,770 cfs with human development, a reduction of 23 percent. More typical summer low flows, described as the summer 95% flow duration, have dropped from 1,760 cfs under unregulated conditions to 1,680 cfs under regulated conditions at the Livingston gage, a reduction of 4.6 percent.

CEA-Related observations in Reach A15 include:

- Passive loss of 2 miles of side channel
- Russian olive colonization in abandoned side channels
- Emergent wetland development in abandoned side channels
- Large corrals that are part of an animal handling facility within 300 feet of the riverbank


Recommended Practices (may include Yellowstone River Recommended Practices--YRRPs) for Reach A15 include:

- Side channel restoration to reactivate 2 miles of passively lost channels
- Russian olive removal (1.2 acres)
- Nutrient management at corrals that are part of an animal handling facility at RM 404
- Consideration of watercraft passage at Big Ditch Diversion Structure
- Consideration of fish passage limitations at Big Ditch Diversion Structure
- Wetland management/restoration due to extent of mapped wetland (150 acres)


The following table summarizes some key CEA results that have been used to describe overall condition and types of human influences affecting the river. The values are specific to this single reach. Blanks indicate that a particular value was not available for this area. This information is consolidated from a large dataset that is presented in more detail in the full reach narrative report.

| | | | | | | |
|---|-------------------------|----------------------------|---|--|---|---|
| Discharge | Undev. | Developed | % Change | "Undeveloped" flows represent conditions prior to significant human development, whereas "developed" flows reflect the current condition of both consumptive and non-consumptive water use. | | |
| 2 Year (cfs) | 31,000 | 29,800 | -3.9% | | | |
| 100 Year (cfs) | 56,600 | 55,900 | -1.2% | | | |
| Bankfull Channel Area (Ac) | 1950 | 1976 | 1995 | 2001 | 1950-2001 | Bankfull channel area is the total footprint of the river inundated at approx. the 2-year flood. |
| | 450.3 | 488.7 | 440.1 | 511.1 | 60.8 | |
| Physical Features | 2011 Length (ft) | % of Bankline | 2001-2011 Change | There are additional types of bank armor such as car bodies and steel retaining walls, but they are relatively minor. | | |
| Rock RipRap | 4,667 | 7.5% | 35 | | | |
| Concrete Riprap | 483 | 0.8% | 0 | | | |
| Flow Deflectors | 0 | 0.0% | 0 | | | |
| Total | 5,150 | 8.3% | 35 | | | |
| Length of Side Channels Blocked (ft) | Pre-1950s | Post-1950s | Numerous side channels have been blocked by small dikes. | | | |
| | 1,617 | 0 | | | | |
| Floodplain Turnover | 1950 - 1976 | 1976 - 2001 | 1950-2001 In-channel riparian encroachment (negative number indicates retreat) | | The rate of floodplain turnover reflects how many acres of land are eroded by the river. Turnover is associated with the creation of riparian habitat. | |
| Total Acres | 141.8 | 120.0 | 4.7 acres | | | |
| Acres/Year | 5.5 | 4.8 | | | | |
| Acres/Year/Valley Mile | 1.1 | 0.9 | | | | |
| Open Bar Area | Point Bars | Bank Attached | Mid-Channel | Total | The type and extent of open sand and gravel bars reflect in-stream habitat conditions that can be important to fish, amphibians, and ground-nesting birds such as least terns. | |
| Change in Area '50 - '01 (Ac) | | | | | | |
| Floodplain Isolation | Acres | % of FP | Floodplain isolation refers to area that historically was flooded, but has become isolated do to flow alterations or physical features such as levees. | | | |
| 5 Year | 27.2 | 25% | | | | |
| 100 Year | 0.0 | 0% | | | | |
| Restricted Migration Area | Acres | % of CMZ | Channel Migration Zone restrictions refer to the area and percent of the CMZ that has been isolated by features such as bank armor, dikes, levees, and transportation embankments. | | | |
| | 122.4 | 8% | | | | |
| Land Use | 1950 | 2011 | 1950 | 2011 | Changes in land use reflect the development of the river corridor through time. The irrigated agricultural are is a sub-set of the mapped agricultural land. | |
| Agricultural Land (Ac) | 2,738.8 | 2,533.8 | Flood (Ac) | 924.9 | 527.9 | |
| Ag. Infrastructure (Ac) | 96.8 | 213.3 | Sprinkler (Ac) | 0.0 | 0.0 | |
| Exurban (Ac) | 0.0 | 2.2 | Pivot (Ac) | 0.0 | 80.5 | |
| Urban (Ac) | 0.0 | 0.0 | | | | |
| Transportation (Ac) | 59.4 | 144.9 | | | | |
| 1950s Riparian Vegetation Converted to a Developed Land Use (ac) | To Irrigated | To Other Use | Total Rip. Converted | % of 1950s Rip. | Changes in the extents of riparian vegetation are influenced by land use changes within the corridor. | |
| | 9.1 | 0.1 | 9.3 | 2.0% | | |
| National Wetlands Inventory | Acres | Acres per Valley Mi | Total Wetland Acres | Wetlands units summarized from National Wetlands Inventory Mapping include Riverine (typically open water sloughs), Emergent (marshes and wet meadows) and Shrub-Scrub (open bar areas with colonizing woody vegetation). | | |
| Riverine | 10.4 | 2.0 | 168.9 | | | |
| Emergent | 131.1 | 25.4 | | | | |
| Scrub/Shrub | 27.4 | 5.3 | | | | |
| Russian Olive (2001) (Appx. 100-yr Floodplain) | Acres | % | Russian olive is considered an invasive species and its presence in the corridor is fairly recent. Its spread can be used as a general indicator of invasive plants within the corridor. | | | |
| | 1.2 | 0.1% | | | | |
| Riparian Forest at low risk of Cowbird Parasitism (Ac/Valley Mile) | 1950 | 1976 | 2001 | Change 1950-2011 | Cowbirds are associated with agricultural and residential development, displacing native bird species by parasitizing their nests. | |
| | 19.9 | 17.5 | 21.2 | 1.2 | | |

PHYSICAL FEATURES MAP (2011)



CHANNEL MIGRATION ZONE MAP



| | | | |
|------------------|--------------------------------------|-----------------------|--------------------|
| County | Stillwater | Upstream River Mile | 400 |
| Classification | PCA: Partially confined anabranching | Downstream River Mile | 392.4 |
| General Location | Park City | Length | 7.60 mi (12.23 km) |

Narrative Summary

Reach A16 is 7.6 miles long and is located just south of Park City. The reach is a Partially Confined Anabranching reach type, indicating some valley wall influences as well as relatively extensive forested islands. The partial geologic confinement within Reach A16 is created by interbedded sandstone and shale. In addition, both low and high alluvial terraces intermittently form the active river corridor margin.

Approximately 9 percent of the bankline in Reach A16 is armored, and the armor is almost entirely rock riprap, some short sections of concrete armor and flow deflectors. The armor is located almost entirely on the northern corridor margin, against terrace margins. Its use is split evenly between protecting agricultural and exurban residential land uses. On the upstream end of the reach, rock armor protects the Italian Ditch Diversion and Canal, which divert water on the north bank of the river at RM 400. Over four miles of floodplain dikes have been mapped in the reach, most of which follow ditches on the north floodplain.

Although there is no evidence that side channels have been intentionally blocked off in Reach A16, there has still been a net loss of over a mile of side channel since 1950. Similar to most reaches in Region A, the loss of side channels has been accompanied by an overall increase in the total channel footprint; since 1950, the bankfull channel area of Reach A16 has increased by 40 acres.

Land use in Reach A16 is almost entirely agricultural, although there are almost 300 acres of urban/exurban development in the mapping footprint. There are corrals that are part of an animal handling facility within 1,000 feet of an abandoned river swale at RM 395. Over a thousand acres under of ground in Reach A16 are under flood irrigation, and about 11 are in pivot. About 150 acres of developed land are in the Channel Migration Zone, and almost 40 acres of that is in urban/exurban development. About 6 percent of the total CMZ is restricted by bank armor and dikes.

There is one pipeline crossing in Reach A16. It crosses under the river at RM 396.7 and consists of a 24 inch crude oil pipeline that is owned by Kinder Morgan Pipelines. This pipeline was horizontally drilled during its installation.

Reach A16 was sampled as part of the avian study. The average species richness in Reach A16 was 8.5, which indicates the average number of species observed during site visits to the reach in cottonwood habitats. The average species richness for all sites evaluated is 8. An average of one cowbird was observed during the field sampling visits. Reach A16 has lost about one half of its riparian forest considered at low risk of cowbird parasitism since 1950. At that time, there were about 12 acres of forest per valley mile considered to be isolated enough from agricultural infrastructure and urban/exurban development to be considered at low risk. By 2011, about 6.6 acres considered low risk remained.

There are over 250 acres of mapped wetland in the reach, with most of that emergent marshes and wet meadows. Many of these wetland areas occupy old river swales on the floodplain north of the river, or abandoned channels in the active corridor.

The reach has extensive Russian olive, with almost 30 acres of mapped footprint in the reach.

A hydrologic evaluation of flow depletions indicates that flow alterations over the last century have been moderate in this reach. The mean annual flood is estimated to have dropped from 16,900 cfs to 15,500 cfs, a drop of about 8 percent. The biggest influence has been on low flows: severe low flows described as 7Q10 (the lowest average 7-day flow anticipated every ten years) for summer months has dropped from an estimated 2,310 cfs to 1,780 cfs with human development, a reduction of 23 percent. More typical summer low flows, described as the summer 95% flow duration, have dropped from 1,760 cfs under unregulated conditions to 1,680 cfs under regulated conditions at the Livingston gage, a reduction of 4.6 percent.

CEA-Related observations in Reach A16 include:

- Passive loss of over a mile of side channel
- Russian olive colonization in abandoned side channels
- Emergent wetland development in abandoned side channels


Recommended Practices (may include Yellowstone River Recommended Practices--YRRPs) for Reach A16 include:

- Diversion structure management at Italian Ditch Diversion RM 400
- Nutrient management at corrals that are part of an animal handling facility at RM 395.
- Russian olive removal (29 acres)
- Wetland management/restoration due to extent of mapped emergent wetland (214 acres emergent, 270 acres total wetland)


The following table summarizes some key CEA results that have been used to describe overall condition and types of human influences affecting the river. The values are specific to this single reach. Blanks indicate that a particular value was not available for this area. This information is consolidated from a large dataset that is presented in more detail in the full reach narrative report.

| | | | | | | |
|---|-------------------------|----------------------------|---|--|---|---|
| Discharge | Undev. | Developed | % Change | "Undeveloped" flows represent conditions prior to significant human development, whereas "developed" flows reflect the current condition of both consumptive and non-consumptive water use. | | |
| 2 Year (cfs) | 32,200 | 30,600 | -5.0% | | | |
| 100 Year (cfs) | 58,600 | 57,600 | -1.7% | | | |
| Bankfull Channel Area (Ac) | 1950 | 1976 | 1995 | 2001 | 1950-2001 | Bankfull channel area is the total footprint of the river inundated at approx. the 2-year flood. |
| | 746.5 | 772.1 | 676.5 | 812.6 | 66.1 | |
| Physical Features | 2011 Length (ft) | % of Bankline | 2001-2011 Change | There are additional types of bank armor such as car bodies and steel retaining walls, but they are relatively minor. | | |
| Rock RipRap | 6,789 | 8.4% | 2,351 | | | |
| Concrete Riprap | 9 | 0.0% | -158 | | | |
| Flow Deflectors | 128 | 0.2% | 128 | | | |
| Total | 6,926 | 8.5% | 2,321 | | | |
| Length of Side Channels Blocked (ft) | Pre-1950s | Post-1950s | Numerous side channels have been blocked by small dikes. | | | |
| | 0 | 0 | | | | |
| Floodplain Turnover | 1950 - 1976 | 1976 - 2001 | 1950-2001 In-channel riparian encroachment (negative number indicates retreat) | The rate of floodplain turnover reflects how many acres of land are eroded by the river. Turnover is associated with the creation of riparian habitat. | | |
| Total Acres | 203.1 | 214.4 | | | | |
| Acres/Year | 7.8 | 8.6 | | | | |
| Acres/Year/Valley Mile | 1.2 | 1.3 | -4.96 acres | | | |
| Open Bar Area | Point Bars | Bank Attached | Mid-Channel | Total | The type and extent of open sand and gravel bars reflect in-stream habitat conditions that can be important to fish, amphibians, and ground-nesting birds such as least terns. | |
| Change in Area '50 - '01 (Ac) | | | | | | |
| Floodplain Isolation | Acres | % of FP | Floodplain isolation refers to area that historically was flooded, but has become isolated do to flow alterations or physical features such as levees. | | | |
| 5 Year | 42.3 | 13% | | | | |
| 100 Year | 0.0 | 0% | | | | |
| Restricted Migration Area | Acres | % of CMZ | Channel Migration Zone restrictions refer to the area and percent of the CMZ that has been isolated by features such as bank armor, dikes, levees, and transportation embankments. | | | |
| | 104.4 | 5% | | | | |
| Land Use | 1950 | 2011 | 1950 | 2011 | Changes in land use reflect the development of the river corridor through time. The irrigated agricultural are is a sub-set of the mapped agricultural land. | |
| Agricultural Land (Ac) | 4,008.9 | 3,532.8 | Flood (Ac) | 1,587.8 | 1,095.2 | |
| Ag. Infrastructure (Ac) | 70.7 | 132.8 | Sprinkler (Ac) | 0.0 | 0.0 | |
| Exurban (Ac) | 0.0 | 268.0 | Pivot (Ac) | 0.0 | 10.6 | |
| Urban (Ac) | 0.0 | 0.0 | | | | |
| Transportation (Ac) | 21.5 | 73.5 | | | | |
| 1950s Riparian Vegetation Converted to a Developed Land Use (ac) | To Irrigated | To Other Use | Total Rip. Converted | % of 1950s Rip. | Changes in the extents of riparian vegetation are influenced by land use changes within the corridor. | |
| | 7.2 | 3.5 | 10.6 | 1.0% | | |
| National Wetlands Inventory | Acres | Acres per Valley Mi | Total Wetland Acres | Wetlands units summarized from National Wetlands Inventory Mapping include Riverine (typically open water sloughs), Emergent (marshes and wet meadows) and Shrub-Scrub (open bar areas with colonizing woody vegetation). | | |
| Riverine | 10.7 | 1.6 | 268.0 | | | |
| Emergent | 214.0 | 32.0 | | | | |
| Scrub/Shrub | 43.3 | 6.5 | | | | |
| Russian Olive (2001) (Appx. 100-yr Floodplain) | Acres | % | Russian olive is considered an invasive species and its presence in the corridor is fairly recent. Its spread can be used as a general indicator of invasive plants within the corridor. | | | |
| | 28.7 | 1.8% | | | | |
| Riparian Forest at low risk of Cowbird Parasitism (Ac/Valley Mile) | 1950 | 1976 | 2001 | Change 1950-2011 | Cowbirds are associated with agricultural and residential development, displacing native bird species by parasitizing their nests. | |
| | 12.1 | 14.5 | 6.6 | -5.5 | | |

PHYSICAL FEATURES MAP (2011)



CHANNEL MIGRATION ZONE MAP



| | | | |
|------------------|-----------------------------|-----------------------|--------------------|
| County | Yellowstone | Upstream River Mile | 392.4 |
| Classification | UA: Unconfined anabranching | Downstream River Mile | 386 |
| General Location | To Laurel | Length | 6.40 mi (10.30 km) |

Narrative Summary

Reach A17 is 7.6 miles long and is located just above Laurel. The reach is classified as Unconfined Anabranching (UA), which is characteristically one of the most dynamic reach types on the river. The river is flowing in the alluvial valley with minimal influences of the valley wall and through numerous forested islands. There are sites in Reach A17 where the river has migrated almost 1,000 feet since 1950.

Approximately 13 percent of the bankline in Reach A17 is armored by rock riprap, concrete riprap and flow deflectors. Between 2001 and 2011 the total length of rock riprap increased by about a half of a mile. At RM 387, a ~750 foot long stretch of flow deflectors on the left bank have been flanked, and by fall 2011 the river had migrated about 120 feet behind the flanked armor. The deflectors are still visible in the channel. In some places such as at RM 389.8, bank armor on both sides of the river narrows the corridor to about one channel width, or 1,000 feet.

Over a mile of side channels in Reach A17 were blocked prior to 1950. Two major channels were blocked on the north side of the river, one at the Buffalo Mirage Fishing Access Site at RM 391.5, and the other at Rm 389.5. These channels, as well as other secondary channels that were passively lost, host fairly dense concentrations of Russian olive. Similar to most reaches in Region A, the loss of side channels has been accompanied by an increase in the total river footprint, indicating that flow concentration into the main river channel has caused it to enlarge. Between 1950 and 2001, the size of the channel increased from 560 acres to 645 acres.

Land use in Reach A17 is primarily agricultural, although there are almost 600 acres of urban/exurban development in the reach as the river approaches the City of Laurel. Since 1950, there has been a reduction in flood irrigated acres of about 550 acres, and an increase in pivot irrigation from 0 acres in 1950 to 284 acres in 2011. A total of 383 acres of developed ground are in the mapped Channel Migration Zone; and about 11 percent of the CMZ has been isolated by physical features protecting those land uses.

At RM 388.5, a headgate diverts water into an old side channel that has been converted to a canal on the north side of the river. About ½ mile downstream, the canal is riprapped where it was recently threatened by rapid northward river migration. At this location, the river has migrated over 800 feet northward since 1950. The main channel of the river now flows along the riprapped canal embankment for about 750 feet.

There are corrals that are part of an animal handling facility within 600 feet of the north riverbank at RM 392.

Side channel loss and channel migration in Reach A17 has resulted in relatively high rates of riparian recruitment. Since 1950, there has been 330 acres of land that experience recruitment of new riparian vegetation. Most of that recruitment was in abandoned channels (200 acres) and about 27 acres of recruitment was direct result of channel migration.

Two ice jams have been recorded in Reach A17, in 1996 and 1997. Both occurred during the month of February, and were reported to have occurred at the Laurel Bridge.

There are over 200 acres of mapped wetland in the reach, with most of that emergent marshes and wet meadows. Many of these wetland areas occupy river swales on the floodplain north of the river, or abandoned channels in the active corridor.

Almost 22 acres of Russian olive has been mapped in the floodplain.

Reach A17 was sampled as part of the avian study. The average species richness in Reach A17 was 7.7, which indicates the average number of species observed during site visits to the reach in cottonwood habitats. The average species richness for all sites evaluated is 8. An average of 0.9 Cowbirds (a bird that parasitizes other bird's nests) were observed in cottonwood habitats during the field sampling visits. Reach A17 has lost about two thirds of its riparian forest considered at low risk of cowbird parasitism since 1950. At that time, there were about 28 acres of forest per valley mile considered to be isolated enough from agricultural infrastructure and urban/exurban development to be considered at low risk. By 2011, about 10 acres per valley mile considered low risk remained.

A total of three Potential Species of Concern (PSOCs) were observed in Reach A17 during the avian study, including the Black and White Warbler, Chimney Swift, and Ovenbird. One Species of Concern (SOC), the Bobolink, was also observed in Reach A17.

A hydrologic evaluation of flow depletions indicates that flow alterations over the last century have been moderate in this reach. The mean annual flood is estimated to have dropped from 16,900 cfs to 15,500 cfs, a drop of about 8 percent. The biggest influence has been on low flows: severe low flows described as 7Q10 (the lowest average 7-day flow anticipated every ten years) for summer months has dropped from an estimated 2,320 cfs to 1,780 cfs with human development, a reduction of 23 percent. More typical summer low flows, described as the summer 95% flow duration, have dropped from 1,760 cfs under unregulated conditions to 1,680 cfs under regulated conditions at the Livingston gage, a reduction of 4.6 percent.

CEA-Related observations in Reach A17 include:

- Flanking of flow deflectors and accelerated erosion behind flanked structures
- Physical blockage of over a mile of side channel
- Russian olive colonization in abandoned side channels

- Emergent wetland development in abandoned side channels
- Ice jamming potentially associated with the Laurel Bridge

Recommended Practices (may include Yellowstone River Recommended Practices--YRRPs) for Reach A17 include:

- Bank armor removal (flanked flow deflectors), RM 387
- Side channel restoration at RM 391.5 and RM 389.5
- Nutrient management associated with corrals that are part of an animal handling facility at RM 392.
- Russian olive removal (22 acres)
- Wetland management/restoration due to extent of mapped wetland (200 acres)
- Irrigation diversion structure management at headgate on side channel at RM 388.5

The following table summarizes some key CEA results that have been used to describe overall condition and types of human influences affecting the river. The values are specific to this single reach. Blanks indicate that a particular value was not available for this area. This information is consolidated from a large dataset that is presented in more detail in the full reach narrative report.

| | | | | | | |
|---|-------------------------|----------------------------|---|--|---|---|
| Discharge | Undev. | Developed | % Change | "Undeveloped" flows represent conditions prior to significant human development, whereas "developed" flows reflect the current condition of both consumptive and non-consumptive water use. | | |
| 2 Year (cfs) | 32,200 | 30,600 | -5.0% | | | |
| 100 Year (cfs) | 58,600 | 57,600 | -1.7% | | | |
| Bankfull Channel Area (Ac) | 1950 | 1976 | 1995 | 2001 | 1950-2001 | Bankfull channel area is the total footprint of the river inundated at approx. the 2-year flood. |
| | 560.0 | 608.9 | 557.5 | 644.6 | 84.6 | |
| Physical Features | 2011 Length (ft) | % of Bankline | 2001-2011 Change | There are additional types of bank armor such as car bodies and steel retaining walls, but they are relatively minor. | | |
| Rock RipRap | 6,184 | 9.1% | 2,584 | | | |
| Concrete Riprap | 2,205 | 3.2% | 0 | | | |
| Flow Deflectors | 671 | 1.0% | -176 | | | |
| Total | 9,060 | 13.3% | 2,407 | | | |
| Length of Side Channels Blocked (ft) | Pre-1950s | Post-1950s | Numerous side channels have been blocked by small dikes. | | | |
| | 7,639 | 0 | | | | |
| Floodplain Turnover | 1950 - 1976 | 1976 - 2001 | 1950-2001 In-channel riparian encroachment (negative number indicates retreat) | The rate of floodplain turnover reflects how many acres of land are eroded by the river. Turnover is associated with the creation of riparian habitat. | | |
| Total Acres | 195.3 | 180.6 | -19.75 acres | | | |
| Acres/Year | 7.5 | 7.2 | | | | |
| Acres/Year/Valley Mile | 1.3 | 1.3 | | | | |
| Open Bar Area | Point Bars | Bank Attached | Mid-Channel | Total | The type and extent of open sand and gravel bars reflect in-stream habitat conditions that can be important to fish, amphibians, and ground-nesting birds such as least terns. | |
| Change in Area '50 - '01 (Ac) | | | | | | |
| Floodplain Isolation | Acres | % of FP | Floodplain isolation refers to area that historically was flooded, but has become isolated do to flow alterations or physical features such as levees. | | | |
| 5 Year | 46.4 | 9% | | | | |
| 100 Year | 89.9 | 7% | | | | |
| Restricted Migration Area | Acres | % of CMZ | Channel Migration Zone restrictions refer to the area and percent of the CMZ that has been isolated by features such as bank armor, dikes, levees, and transportation embankments. | | | |
| | 245.6 | 11% | | | | |
| Land Use | 1950 | 2011 | 1950 | 2011 | Changes in land use reflect the development of the river corridor through time. The irrigated agricultural are is a sub-set of the mapped agricultural land. | |
| Agricultural Land (Ac) | 4,530.2 | 4,110.3 | Flood (Ac) | 1,927.0 | 1,384.1 | |
| Ag. Infrastructure (Ac) | 68.6 | 118.5 | Sprinkler (Ac) | 0.0 | 0.0 | |
| Exurban (Ac) | 59.1 | 292.3 | Pivot (Ac) | 0.0 | 283.8 | |
| Urban (Ac) | 95.4 | 203.9 | | | | |
| Transportation (Ac) | 50.2 | 50.2 | | | | |
| 1950s Riparian Vegetation Converted to a Developed Land Use (ac) | To Irrigated | To Other Use | Total Rip. Converted | % of 1950s Rip. | Changes in the extents of riparian vegetation are influenced by land use changes within the corridor. | |
| | 6.0 | 0.8 | 6.8 | 1.0% | | |
| National Wetlands Inventory | Acres | Acres per Valley Mi | Total Wetland Acres | Wetlands units summarized from National Wetlands Inventory Mapping include Riverine (typically open water sloughs), Emergent (marshes and wet meadows) and Shrub-Scrub (open bar areas with colonizing woody vegetation). | | |
| Riverine | 9.4 | 1.6 | 226.2 | | | |
| Emergent | 203.4 | 35.6 | | | | |
| Scrub/Shrub | 13.4 | 2.3 | | | | |
| Russian Olive (2001) (Appx. 100-yr Floodplain) | Acres | % | Russian olive is considered an invasive species and its presence in the corridor is fairly recent. Its spread can be used as a general indicator of invasive plants within the corridor. | | | |
| | 21.8 | 6.7% | | | | |
| Riparian Forest at low risk of Cowbird Parasitism (Ac/Valley Mile) | 1950 | 1976 | 2001 | Change 1950-2011 | Cowbirds are associated with agricultural and residential development, displacing native bird species by parasitizing their nests. | |
| | 27.7 | 64.2 | 9.7 | -18.0 | | |