

County	Yellowstone	Upstream River Mile	368.3
Classification	PCB: Partially confined braided	Downstream River Mile	362.2
General Location	Billings	Length	6.10 mi (9.82 km)

Narrative Summary

Reach B2 is 6.1 miles long and located in Billings. The reach extends from the rimrock bluffs south of town, under the I-90 Bridge, to the refinery area at Lockwood. It is a Partially Confined Braided (PCB) reach type indicating some influence of the bluff line on the river coupled by extensive open gravel bars and low flow channels. Reach B2 is extensively urbanized, with floodplain dikes, industrial and urban/exurban development, pipeline crossings, and bridges throughout the reach. Flow alterations in this reach have been substantial; the mean annual flood has dropped an estimated 17 percent due to human influences, and summer low flows have dropped by 42 percent.

In total there are 21,700 feet of bank armor in Reach B2, which equates to 4.1 miles of bank armor in a 6 mile long reach of river. Concrete riprap is the most prevalent type of armor, with about three miles present in 2011. There is almost a mile of rock riprap and a few flow deflectors. There are also over three miles of floodplain dikes mapped in the reach.

Since 1950, 6,566 feet of side channels have been blocked by dikes. These blocked side channels are in highly urbanized areas upstream of the I-90 Bridge and at the water treatment plant downstream.

The primary land use in the reach is urban/exurban development. A total of 620 acres of the historic 100-year floodplain has become isolated from the river, which is 41 percent of the total 100-year floodplain footprint. Most of the 100-year floodplain isolation is due to the Interstate Highway Embankment. Approximately 21 percent of the Channel Migration Zone has become restricted due to physical features, most of which are riprap installed to protect urban/industrial land uses.

A total of three ice jams have been recorded in Reach B2. One of these jams occurred in February of 1996, and the other two in January of 1997. They all resulted in flooding and the January 3 1997 jam caused some evacuations. The jams were reported as forming upstream of the I-90 Bridge.

There are numerous pipeline crossings in Reach B2. At RM 367 two pipelines cross under the river. One is a crude oil pipeline owned by Beartooth Pipeline that is HDD (Horizontal Directionally Drilled). The other is a petroleum product pipeline owned by Phillips 66 that as of Fall 2012 was trenched, and according to the addendum to the Yellowstone River Pipeline Risk Assessment, had 4 to 10 feet of cover. Further downstream, there are seven pipelines listed in the Pipeline Risk Assessment Report at RM 365. Several of these pipelines are trenched as a bundle, with a reported minimum of two feet of cover.

About 25 acres of Russian olive have been mapped in Reach B2.

Reach B2 was sampled as part of the fisheries study. A total of 31 fish species were sampled in the reach and one of those species was Sauger, which has been identified by the Montana Natural Heritage Program as a Species of Concern (SOC).

Reach B2 was sampled as part of the avian study. The average species richness in Reach B2 was 7.0, which indicates the average number of species observed during site visits to the reach in cottonwood habitats. The average species richness for sites evaluated is 8. Two bird species identified by the Montana Natural Heritage Program as Potential Species of Concern (PSOC) were also found, the Ovenbird and the Plumbeous Vireo.

A hydrologic evaluation of flow depletions indicates that flow alterations over the last century have been substantial in this reach. The mean annual flood is estimated to have dropped from 23,700 cfs to 19,700 cfs, a drop of about 17 percent. Low flows have also been impacted; severe low flows described as 7Q10 (the lowest average 7-day flow anticipated every ten years) for summer months has dropped from an estimated 2,910 cfs to 2,000 cfs with human development, a reduction of 31 percent. More typical summer low flows, described as the summer 95% flow duration, have dropped from 3,836 cfs under unregulated conditions to 2,227 cfs under regulated conditions at the Billings gage, a reduction of 42 percent.

CEA-Related observations in Reach B2 include:

- Extensive armoring with CMZ encroachment


Recommended Practices (may include Yellowstone River Recommended Practices--YRRPs) for Reach B2 include:

- Pipeline crossing management
- Russian olive removal

The following table summarizes some key CEA results that have been used to describe overall condition and types of human influences affecting the river. The values are specific to this single reach. Blanks indicate that a particular value was not available for this area. This information is consolidated from a large dataset that is presented in more detail in the full reach narrative report.

Discharge	Undev.	Developed	% Change	"Undeveloped" flows represent conditions prior to significant human development, whereas "developed" flows reflect the current condition of both consumptive and non-consumptive water use.		
2 Year (cfs)	44,200	39,800	-10.0%			
100 Year (cfs)	78,600	76,000	-3.3%			
Bankfull Channel Area (Ac)	1950	1976	1995	2001	1950-2001	Bankfull channel area is the total footprint of the river inundated at approx. the 2-year flood.
	517.8	536.9	501.3	534.2	16.4	
Physical Features	2011 Length (ft)	% of Bankline	2001-2011 Change	There are additional types of bank armor such as car bodies and steel retaining walls, but they are relatively minor.		
Rock Riprap	4,329	6.7%	828			
Concrete Riprap	17,283	26.8%	0			
Flow Deflectors	91	0.1%	91			
Total	21,702	33.7%	918			
Length of Side Channels Blocked (ft)	Pre-1950s	Post-1950s	Numerous side channels have been blocked by small dikes.			
	0	6,566				
Floodplain Turnover	1950 - 1976	1976 - 2001	1950-2001 In-channel riparian encroachment (negative number indicates retreat)	The rate of floodplain turnover reflects how many acres of land are eroded by the river. Turnover is associated with the creation of riparian habitat.		
Total Acres	136.5	88.0	-37.22 acres			
Acres/Year	5.3	3.5				
Acres/Year/Valley Mile	0.9	0.6				
Open Bar Area	Point Bars	Bank Attached	Mid-Channel	Total	The type and extent of open sand and gravel bars reflect in-stream habitat conditions that can be important to fish, amphibians, and ground-nesting birds such as least terns.	
Change in Area '50 - '01 (Ac)						
Floodplain Isolation	Acres	% of FP	Floodplain isolation refers to area that historically was flooded, but has become isolated do to flow alterations or physical features such as levees.			
5 Year	58.1	15%				
100 Year	620.1	41%				
Restricted Migration Area	Acres	% of CMZ	Channel Migration Zone restrictions refer to the area and percent of the CMZ that has been isolated by features such as bank armor, dikes, levees, and transportation embankments.			
	255.5	21%				
Land Use	1950	2011	1950	2011	Changes in land use reflect the development of the river corridor through time. The irrigated agricultural are is a sub-set of the mapped agricultural land.	
Agricultural Land (Ac)	2,457.5	1,071.5	Flood (Ac)	469.3	0.0	
Ag. Infrastructure (Ac)	33.0	17.2	Sprinkler (Ac)	0.0	5.5	
Exurban (Ac)	318.3	0.0	Pivot (Ac)	0.0	0.0	
Urban (Ac)	760.2	2,495.1				
Transportation (Ac)	46.0	127.8				
1950s Riparian Vegetation Converted to a Developed Land Use (ac)	To Irrigated	To Other Use	Total Rip. Converted	% of 1950s Rip.	Changes in the extents of riparian vegetation are influenced by land use changes within the corridor.	
	0.0	317.3	317.3	51.0%		
National Wetlands Inventory	Acres	Acres per Valley Mi	Total Wetland Acres	Wetlands units summarized from National Wetlands Inventory Mapping include Riverine (typically open water sloughs), Emergent (marshes and wet meadows) and Shrub-Scrub (open bar areas with colonizing woody vegetation).		
Riverine	44.5	8.0	75.7			
Emergent	19.6	3.5				
Scrub/Shrub	11.6	2.1				
Russian Olive (2001) (Appx. 100-yr Floodplain)	Acres	%	Russian olive is considered an invasive species and its presence in the corridor is fairly recent. Its spread can be used as a general indicator of invasive plants within the corridor.			
	24.6	3.2%				
Riparian Forest at low risk of Cowbird Parasitism (Ac/Valley Mile)	1950	1976	2001	Change 1950-2011	Cowbirds are associated with agricultural and residential development, displacing native bird species by parasitizing their nests.	
	5.0	1.9	4.0	-1.0		

PHYSICAL FEATURES MAP (2011)



CHANNEL MIGRATION ZONE MAP

