

County	Yellowstone	Upstream River Mile	322.7
Classification	UA: Unconfined anabranching	Downstream River Mile	318
General Location	Reed Creek	Length	4.70 mi (7.56 km)

### Narrative Summary

Reach B9 is located in lower Yellowstone County near Reed Creek. The Reach is 4.7 miles long and is an Unconfined Anabranching (UA) reach type, indicating the presence of extensive forested islands with little valley wall influence on the main channel. This reach type is typically the most dynamic in the system due to a lack of confinement and extent of side channels.

About 7,300 feet of streambank are armored by rock riprap, which is about 15 percent of the total bankline. Most of the bank armor in the reach is protecting the rail line on the south side of the river, and most of it is located along the edge of a section of bluff line. Another section of armor is protecting a major power line crossing on the north bank at RM 321. Currently, two towers on the crossing are right on the edge of the river.

One side channel that is about 8,000 feet long at RM 321.5L was blocked prior to 1950. The lower end of this old channel still holds open water, but the upstream end has been graded into fields and also supports two major power line towers.

Land uses related to both irrigation and the railroad have encroached into the Channel Migration Zone (CMZ) in Reach B9. Overall, land uses in the reach are primarily agricultural, with about 508 acres of flood irrigated land mapped as of 2011. About half of that irrigated acreage is within the CMZ. There are 384 acres under pivot, about 75 of which are within the CMZ. The railroad has encroached into 101 acres of the CMZ and is primarily responsible for its isolation. In total, just under 10 percent of the CMZ has been restricted due to bank armor, and 7.3 percent of the restriction is due to the railroad, while 2.4 percent is associated with the protection of irrigated lands.

The modern 5-year floodplain contains about 76 acres of flood-irrigated ground, and 64 acres of ground under pivot.

Waco-Custer Diversion Dam is located at RM 320. The Waco-Custer ditch company was formed in the early 1900's, and the diversion dam was constructed shortly thereafter (<http://www.fws.gov/YellowstoneRiverCoordinator/Waco-custer.html>). The Waco-Custer diversion supports approximately 4,300 acres of irrigation, with a diversion capacity of 125 cfs. The structure is located approximately eight miles west of Custer, at River Mile 320. At the diversion, the Yellowstone River flows through two main channels, and the structure itself blocks only the right channel. The structure feeds the Waco-Custer Canal, which flows on the south floodplain surface of the Yellowstone River.

Migration rates in several locations in Reach B9 have exceeded an average of 10 feet per year since the mid-1950s. At Rm 322, the river migrated almost 200 feet between 2001 and 2011, which is double that average rate of 10 feet per year. That rapid recent migration has been through irrigated fields on the south side of the river. Lateral migration of the river has promoted extensive recruitment of new woody riparian habitat. Since the 1950s there has been about 210 acres of riparian recruitment in the reach, most of which was riparian colonization of old 1950's channel area. Additionally, there are 213 mapped wetlands in the reach, including 105 acres of emergent wetland types such as wet meadows and marsh. The reach contains about 53 wetland acres per valley mile, which is a relatively high value for the Yellowstone River.

Reach B9 has had a major loss of forest area that is considered at low risk of cowbird parasitism. In 1950, there were about 48 acres per valley mile of such forest, and that had been reduced by 2001 to 21 acres per valley mile.

A hydrologic evaluation of flow depletions indicates that flow alterations over the last century have been major in this reach. The mean annual flood is estimated to have dropped from 30,200 cfs to 24,500 cfs, a drop of about 19 percent. The 2-year flood, which strongly influences overall channel form, has dropped by 11 percent. Low flows have also been impacted; severe low flows described as 7Q10 (the lowest average 7-day flow anticipated every ten years) for summer months has dropped from an estimated 3,060 cfs to 2,080 cfs with human development, a reduction of 32 percent. More typical summer low flows, described as the summer 95% flow duration, have dropped from 3,846 cfs under unregulated conditions to 2,227 cfs under regulated conditions at the Billings gage, a reduction of 42 percent.

About 23 percent of the 5-year floodplain has become isolated in Reach B9, and the vast majority of this isolation is on the south side of the river at RM 321 where the rail line has isolated an historic side channel. Much of that 5-year floodplain isolation is due to transportation infrastructure on the south side of the river. This isolated floodplain area still holds open water in a distinct swale.

CEA-Related observations in Reach B9 include:

- Blockage of one side channel at RM 321.5 sometime prior to 1950
- Railroad isolation of major channel remnant that supports open water.

Recommended Practices (may include Yellowstone River Recommended Practices--YRRPs) for Reach B9 include:

- Side channel reactivation at RM 321.5—may be difficult due to power line
- CMZ management due to ~10 percent restriction of CMZ
- Russian olive removal
- Floodplain reconnection where active rail line has isolated historic channel remnant at RM 321R.
- Fish passage Practice at Waco Custer Diversion Dam (not complete blockage)
- Watercraft passage Practice at Waco Custer Diversion Dam (side channel passage exists)
- Irrigation Infrastructure management at Waco Custer Diversion Dam.

The following table summarizes some key CEA results that have been used to describe overall condition and types of human influences affecting the river. The values are specific to this single reach. Blanks indicate that a particular value was not available for this area. This information is consolidated from a large dataset that is presented in more detail in the full reach narrative report.

<b>Discharge</b>	<b>Undev.</b>	<b>Developed</b>	<b>% Change</b>	<b>"Undeveloped" flows represent conditions prior to significant human development, whereas "developed" flows reflect the current condition of both consumptive and non-consumptive water use.</b>		
2 Year (cfs)	55,500	49,400	-11.0%			
100 Year (cfs)	97,200	93,600	-3.7%			
<b>Bankfull Channel Area (Ac)</b>	<b>1950</b>	<b>1976</b>	<b>1995</b>	<b>2001</b>	<b>1950-2001</b>	<b>Bankfull channel area is the total footprint of the river inundated at approx. the 2-year flood.</b>
	485.8	524.8	515.2	539.2	53.5	
<b>Physical Features</b>	<b>2011 Length (ft)</b>	<b>% of Bankline</b>	<b>2001-2011 Change</b>	<b>There are additional types of bank armor such as car bodies and steel retaining walls, but they are relatively minor.</b>		
Rock RipRap	7,304	14.9%	0			
Concrete Riprap	0	0.0%	0			
Flow Deflectors	89	0.2%	0			
<b>Total</b>	<b>7,393</b>	<b>15.1%</b>	<b>0</b>			
<b>Length of Side Channels Blocked (ft)</b>	<b>Pre-1950s</b>	<b>Post-1950s</b>	<b>Numerous side channels have been blocked by small dikes.</b>			
	7,943	0				
<b>Floodplain Turnover</b>	<b>1950 - 1976</b>	<b>1976 - 2001</b>	<b>1950-2001 In-channel riparian encroachment (negative number indicates retreat)</b>		<b>The rate of floodplain turnover reflects how many acres of land are eroded by the river. Turnover is associated with the creation of riparian habitat.</b>	
Total Acres	166.0	162.6	6.4 acres			
Acres/Year	6.4	6.5				
Acres/Year/Valley Mile	1.6	1.7				
<b>Open Bar Area</b>	<b>Point Bars</b>	<b>Bank Attached</b>	<b>Mid-Channel</b>	<b>Total</b>	<b>The type and extent of open sand and gravel bars reflect in-stream habitat conditions that can be important to fish, amphibians, and ground-nesting birds such as least terns.</b>	
<b>Change in Area '50 - '01 (Ac)</b>						
<b>Floodplain Isolation</b>	<b>Acres</b>	<b>% of FP</b>	<b>Floodplain isolation refers to area that historically was flooded, but has become isolated do to flow alterations or physical features such as levees.</b>			
5 Year	175.0	23%				
100 Year	0.0	0%				
<b>Restricted Migration Area</b>	<b>Acres</b>	<b>% of CMZ</b>	<b>Channel Migration Zone restrictions refer to the area and percent of the CMZ that has been isolated by features such as bank armor, dikes, levees, and transportation embankments.</b>			
	168.5	10%				
<b>Land Use</b>	<b>1950</b>	<b>2011</b>	<b>1950</b>	<b>2011</b>	<b>Changes in land use reflect the development of the river corridor through time. The irrigated agricultural are is a sub-set of the mapped agricultural land.</b>	
Agricultural Land (Ac)	2,906.3	2,697.0	Flood (Ac)	656.7	507.8	
Ag. Infrastructure (Ac)	12.0	62.4	Sprinkler (Ac)	0.0	0.0	
Exurban (Ac)	0.6	0.6	Pivot (Ac)	0.0	384.1	
Urban (Ac)	0.0	0.0				
Transportation (Ac)	61.4	153.2				
<b>1950s Riparian Vegetation Converted to a Developed Land Use (ac)</b>	<b>To Irrigated</b>	<b>To Other Use</b>	<b>Total Rip. Converted</b>	<b>% of 1950s Rip.</b>	<b>Changes in the extents of riparian vegetation are influenced by land use changes within the corridor.</b>	
	4.9	0.5	5.4	1.0%		
<b>National Wetlands Inventory</b>	<b>Acres</b>	<b>Acres per Valley Mi</b>	<b>Total Wetland Acres</b>		<b>Wetlands units summarized from National Wetlands Inventory Mapping include Riverine (typically open water sloughs), Emergent (marshes and wet meadows) and Shrub-Scrub (open bar areas with colonizing woody vegetation).</b>	
Riverine	24.3	6.2	212.5			
Emergent	104.6	26.9				
Scrub/Shrub	83.6	21.5				
<b>Russian Olive (2001) (Appx. 100-yr Floodplain)</b>	<b>Acres</b>	<b>%</b>	<b>Russian olive is considered an invasive species and its presence in the corridor is fairly recent. Its spread can be used as a general indicator of invasive plants within the corridor.</b>			
	5.9	0.3%				
<b>Riparian Forest at low risk of Cowbird Parasitism (Ac/Valley Mile)</b>	<b>1950</b>	<b>1976</b>	<b>2001</b>	<b>Change 1950-2011</b>	<b>Cowbirds are associated with agricultural and residential development, displacing native bird species by parasitizing their nests.</b>	
	47.7	28.0	21.0	-26.7		



## CHANNEL MIGRATION ZONE MAP

