

<b>County</b>	Mckenzie	<b>Upstream River Mile</b>	7.5
<b>Classification</b>	US/I: Unconfined straight/islands	<b>Downstream River Mile</b>	0
<b>General Location</b>	To Missouri River	<b>Length</b>	7.50 mi (12.07 km)

## Narrative Summary

Reach D16 is the lowermost reach of the Yellowstone River, extending 7.5 miles to the confluence with the Missouri River. It is a unique reach type, referred to as Unconfined Straight (US), and it has numerous forested islands that have developed since the 1950s.

Reach D16 has only a few hundred feet of rock riprap along its 7.5 mile length, and all of that was built since 2001. No side channels have been blocked.

The most striking change in Reach D16 since 1950 is the encroachment of riparian vegetation onto old sand bars. Between 1950 and 2001, the size of the channel has dropped by 550 acres, and there has been 472 acres of riparian encroachment into old channel areas. Much of this encroachment converted open sand bars into forested islands. There has been a loss of over 150 acres of sand bar since 1950. This change has resulted in a conversion of almost 7 miles low flow channels around gravel bars to anabranching side channels around islands.

Land use in the reach is dominated by flood irrigation. The extent of flood irrigated lands increased from 4,600 acres in 1950 to about 8,500 acres in 2011. The floodplain is very flat and broad in this lowermost portion of the Yellowstone River valley, and as a result, floodplain development for agriculture has substantially altered floodplain access. About 29 percent of the 100-year floodplain has become isolated from the river, and a fraction of this (1.6 percent) has been attributed to flow alterations, whereas 27 percent has been associated with agricultural features on the floodplain such as roads and ditches. There are about 480 acres of flood irrigated land within the Channel Migration Zone of Reach D16.

Land use mapping shows several drill pads in the lower portion of the reach that are within several thousand feet of the river. There are four drill pads on a narrow strip of land at the mouth that lies between the Yellowstone and Missouri Rivers.

Reach D16 has a notably high concentration of mapped wetlands. There are about 580 acres of mapped wetland in the reach, which translates to about 80 acres per valley mile. Along the rest of the river, wetland densities rarely exceed 50 acres per valley mile. Reach D16 only has 3.5 acres of mapped Russian olive, which is a relatively low density for reaches below Billings.

Because of the riparian encroachment, Reach D16 has seen an increase in the area of riparian forest considered at low risk of cowbird parasitism; in 1950 there were about 250 acres of such forest per valley mile, and in 2001 there were 308 acres per valley mile.

The changes in Reach D16 are due in part to major flow alterations in the reach. The 2-year discharge, which is considered to have a large influence on channel size, has been reduced by 22 percent due to human development.

CEA-Related observations in Reach D16 include:

- Extensive riparian encroachment with flow alterations
- Conversion of open sand bars to forested islands


Recommended Practices (may include Yellowstone River Recommended Practices--YRRPs) for Reach D16 include:

- Drill pad considerations
- Riparian protections

The following table summarizes some key CEA results that have been used to describe overall condition and types of human influences affecting the river. The values are specific to this single reach. Blanks indicate that a particular value was not available for this area. This information is consolidated from a large dataset that is presented in more detail in the full reach narrative report.

<b>Discharge</b>	<b>Undev.</b>	<b>Developed</b>	<b>% Change</b>	<b>"Undeveloped" flows represent conditions prior to significant human development, whereas "developed" flows reflect the current condition of both consumptive and non-consumptive water use.</b>		
2 Year (cfs)	69,900	54,300	-22.3%			
100 Year (cfs)	143,000	134,000	-6.3%			
<b>Bankfull Channel Area (Ac)</b>	<b>1950</b>	<b>1976</b>	<b>1995</b>	<b>2001</b>	<b>1950-2001</b>	<b>Bankfull channel area is the total footprint of the river inundated at approx. the 2-year flood.</b>
	1,515.1		1,157.3	960.1	-555.0	
<b>Physical Features</b>	<b>2011 Length (ft)</b>	<b>% of Bankline</b>	<b>2001-2011 Change</b>	<b>There are additional types of bank armor such as car bodies and steel retaining walls, but they are relatively minor.</b>		
Rock RipRap	266	0.3%	266			
Concrete Riprap	0	0.0%	0			
Flow Deflectors	0	0.0%	0			
<b>Total</b>	<b>266</b>	<b>0.3%</b>	<b>266</b>			
<b>Length of Side Channels Blocked (ft)</b>	<b>Pre-1950s</b>	<b>Post-1950s</b>	<b>Numerous side channels have been blocked by small dikes.</b>			
	0	0				
<b>Floodplain Turnover</b>	<b>1950 - 1976</b>	<b>1976 - 2001</b>	<b>1950-2001 In-channel riparian encroachment (negative number indicates retreat)</b>		<b>The rate of floodplain turnover reflects how many acres of land are eroded by the river. Turnover is associated with the creation of riparian habitat.</b>	
Total Acres Acres/Year Acres/Year/Valley Mile			472.19 acres			
<b>Open Bar Area</b>	<b>Point Bars</b>	<b>Bank Attached</b>	<b>Mid-Channel</b>	<b>Total</b>	<b>The type and extent of open sand and gravel bars reflect in-stream habitat conditions that can be important to fish, amphibians, and ground-nesting birds such as least terns.</b>	
Change in Area '50 - '01 (Ac)	10.3	45.8	-208.4	-152.3		
<b>Floodplain Isolation</b>	<b>Acres</b>	<b>% of FP</b>	<b>Floodplain isolation refers to area that historically was flooded, but has become isolated do to flow alterations or physical features such as levees.</b>			
5 Year	105.9	31%				
100 Year	390.4	29%				
<b>Restricted Migration Area</b>	<b>Acres</b>	<b>% of CMZ</b>	<b>Channel Migration Zone restrictions refer to the area and percent of the CMZ that has been isolated by features such as bank armor, dikes, levees, and transportation embankments.</b>			
<b>Land Use</b>	<b>1950</b>	<b>2011</b>	<b>1950</b>	<b>2011</b>	<b>Changes in land use reflect the development of the river corridor through time. The irrigated agricultural are is a sub-set of the mapped agricultural land.</b>	
Agricultural Land (Ac)	10,472.2	14,362.1	Flood (Ac)	4,631.0	8,492.4	
Ag. Infrastructure (Ac)	87.1	270.2	Sprinkler (Ac)	0.0	0.0	
Exurban (Ac)	0.0	63.7	Pivot (Ac)	0.0	0.0	
Urban (Ac)	0.0	0.0				
Transportation (Ac)	0.0	17.9				
<b>1950s Riparian Vegetation Converted to a Developed Land Use (ac)</b>	<b>To Irrigated</b>	<b>To Other Use</b>	<b>Total Rip. Converted</b>	<b>% of 1950s Rip.</b>	<b>Changes in the extents of riparian vegetation are influenced by land use changes within the corridor.</b>	
<b>National Wetlands Inventory</b>	<b>Acres</b>	<b>Acres per Valley Mi</b>	<b>Total Wetland Acres</b>	<b>Wetlands units summarized from National Wetlands Inventory Mapping include Riverine (typically open water sloughs), Emergent (marshes and wet meadows) and Shrub-Scrub (open bar areas with colonizing woody vegetation).</b>		
Riverine	25.3	3.6	558.4			
Emergent	254.9	36.2				
Scrub/Shrub	278.2	39.5				
<b>Russian Olive (2001) (Appx. 100-yr Floodplain)</b>	<b>Acres</b>	<b>%</b>	<b>Russian olive is considered an invasive species and its presence in the corridor is fairly recent. Its spread can be used as a general indicator of invasive plants within the corridor.</b>			
	3.5	0.1%				
<b>Riparian Forest at low risk of Cowbird Parasitism (Ac/Valley Mile)</b>	<b>1950</b>	<b>1976</b>	<b>2001</b>	<b>Change 1950-2011</b>	<b>Cowbirds are associated with agricultural and residential development, displacing native bird species by parasitizing their nests.</b>	
	230.3		307.9	77.6		

## PHYSICAL FEATURES MAP (2011)



CHANNEL MIGRATION ZONE MAP

