




Land Cover

Summarized by: **Meagher** (County)



20%
(305,579
Acres)

Shrubland, Steppe and Savanna Systems Sagebrush Steppe

Big Sagebrush Steppe

This widespread ecological system occurs throughout much of central Montana, and north and east onto the western fringe of the Great Plains. In central Montana, where this system occurs on both glaciated and non-glaciated landscapes, it differs slightly, with more summer rain than winter precipitation and more precipitation annually. Throughout its distribution, soils are typically deep and non-saline, often with a microphytic crust. This shrub-steppe is dominated by perennial grasses and forbs with greater than 25% cover. Overall shrub cover is less than 10 percent. In Montana and Wyoming, stands are more mesic, with more biomass of grass, and have less shrub diversity than stands farther to the west, and 50 to 90% of the occurrences are dominated by Wyoming big sagebrush with western wheatgrass (*Pascopyrum smithii*). Japanese brome (*Bromus japonicus*) and cheatgrass (*Bromus tectorum*) are indicators of disturbance, but cheatgrass is typically not as abundant as in the Intermountain West, possibly due to a colder climate. The natural fire regime of this ecological system maintains a patchy distribution of shrubs, preserving the steppe character. Shrubs may increase following heavy grazing and/or with fire suppression. In central and eastern Montana, complexes of prairie dog towns are common in this ecological system.



16%
(240,994
Acres)

Shrubland, Steppe and Savanna Systems

Sagebrush Steppe

Montane Sagebrush Steppe

This system dominates the montane and subalpine landscape of southwestern Montana from valley bottoms to subalpine ridges and is found as far north as Glacier National Park. It can also be seen in the island mountain ranges of the north-central and south-central portions of the state. It primarily occurs on deep-soiled to stony flats, ridges, nearly flat ridgetops, and mountain slopes. In general, this system occurs in areas of gentle topography, fine soils, subsurface moisture or mesic conditions, within zones of higher precipitation and areas of snow accumulation. It occurs on all slopes and aspects, variable substrates and all soil types. The shrub component of this system is generally dominated by mountain big sagebrush (*Artemisia tridentata* ssp. *vaseyana*). Other co-dominant shrubs include silver sagebrush (*Artemisia cana* ssp. *viscidula*), subalpine big sagebrush (*Artemisia tridentata* ssp. *spiciformis*), three tip sagebrush (*Artemisia tripartita* ssp. *tripartita*) and antelope bitterbrush (*Purshia tridentata*). Little sagebrush (*Artemisia arbuscula* ssp. *arbuscula*) shrublands are only found in southwestern Montana on sites with a perched water table. Wyoming big sagebrush (*Artemisia tridentata* ssp. *wyomingensis*) sites may be included within this system if occurrences are at montane elevations, and are associated with montane graminoids such as Idaho fescue (*Festuca idahoensis*), spike fescue (*Leucopoa kingii*), or poverty oatgrass (*Danthonia intermedia*). In areas where sage has been eliminated by human activities like burning, disking or poisoning, other shrubs may be dominant, especially rubber rabbitbrush (*Ericameria nauseosa*), and green rabbitbrush (*Chrysothamnus viscidiflorus*). Because of the mesic site conditions, most occurrences support a diverse herbaceous undergrowth of grasses and forbs. Shrub canopy cover is extremely variable, ranging from 10 percent to as high as 40 or 50 percent.



14%
(221,958
Acres)

Forest and Woodland Systems

Conifer-dominated forest and woodland (xeric-mesic)

Rocky Mountain Montane Douglas-fir Forest and Woodland

In Montana, this ecological system occurs on the east side of the Continental Divide, north to about the McDonald Pass area, and along the Rocky Mountain Front. This system is associated with a dry to submesic continental climate regime with annual precipitation ranging from 51 to 102 centimeters (20-40 inches), with a maximum in winter or late spring. Winter snowpacks typically melt off in early spring at lower elevations. Elevations range from valley bottoms to 1,980 meters (6500 feet) in northern Montana and up to 2,286 meters (7500 feet) on warm aspects in southern Montana. It occurs on north-facing aspects in most areas, and south-facing aspects at higher elevations. This is a Douglas-fir (*Pseudotsuga menziesii*) dominated system without any maritime floristic composition. Fire disturbance intervals are as infrequent as 500 years, and as a result, individual trees and forests can attain great age on some sites (500 to 1,500 years). In Montana, this system occurs from lower montane to lower subalpine environments and is prevalent on calcareous substrates. Common understory shrubs include common ninebark (*Physocarpus malvaceus*), common juniper (*Juniperus communis*), Rocky Mountain juniper (*Juniperus scopulorum*), birch-leaf spiraea (*Spiraea betulifolia*), snowberry (*Symphoricarpos* species), creeping Oregon grape (*Mahonia repens*) and Canadian buffaloberry (*Shepherdia canadensis*). The Douglas-fir/pinegrass (*Calamagrostis rubescens*) type is the most ubiquitous association found within this system in Montana.



11%
(173,606
Acres)

Grassland Systems

Montane Grassland

Rocky Mountain Lower Montane, Foothill, and Valley Grassland

This grassland system of the northern Rocky Mountains is found at lower montane to foothill elevations in mountains and valleys throughout Montana. These grasslands are floristically similar to Big Sagebrush Steppe but are defined by shorter summers, colder winters, and young soils derived from recent glacial and alluvial material. They are found at elevations from 548 - 1,650 meters (1,800-5,413 feet). In the lower montane zone, they range from small meadows to large open parks surrounded by conifers; below the lower treeline, they occur as extensive foothill and valley grasslands. Soils are relatively deep, fine-textured, often with coarse fragments, and non-saline. Microphytic crust may be present in high-quality occurrences. This system is typified by cool-season perennial bunch grasses and forbs (>25%) cover, with a sparse shrub cover (<10%). Rough fescue (*Festuca campestris*) is dominant in the northwestern portion of the state and Idaho fescue (*Festuca idahoensis*) is dominant or co-dominant throughout the range of the system. Bluebunch wheatgrass (*Pseudoroegneria spicata*) occurs as a co-dominant throughout the range as well, especially on xeric sites. Western wheatgrass (*Pascopyrum smithii*) is consistently present, often with appreciable coverage (>10%) in lower elevation occurrences in western Montana and virtually always present, with relatively high coverages (>25%), on the edge of the Northwestern Great Plains region. Species diversity ranges from a high of more than 50 per 400 square meter plot on mesic sites to 15 (or fewer) on xeric and disturbed sites. Most occurrences have at least 25 vascular species present. Farmland conversion, noxious species invasion, fire suppression, heavy grazing and oil and gas development are major threats to this system.



11%
(168,887
Acres)

Forest and Woodland Systems Conifer-dominated forest and woodland (xeric-mesic)

Rocky Mountain Lodgepole Pine Forest

This forested system is widespread in upper montane to subalpine zones of the Montana Rocky Mountains, and east into island ranges of north-central Montana and the Bighorn and Beartooth ranges of south-central Montana. These are montane to subalpine forests where the dominance of lodgepole pine (*Pinus contorta*) is related to fire history and topographic conditions. In Montana, elevation ranges from 975 to 2,743 meters (3,200-9000 feet). These forests occur on flats to slopes of all degrees and aspect, as well as valley bottoms. Fire is frequent, and stand-replacing fires are common. Following stand-replacing fires, lodgepole pinewill rapidly colonize and develop into dense, even-aged stands. Most forests in this ecological system occur as early- to mid-successional forests persisting for 50-200 years on warmer, lower elevation forests, and 150-400 years in subalpine forests. They generally occur on dry to intermediate sites with a wide seasonal range of temperatures and long precipitation-free periods in summer. Snowfall is heavy and supplies the major source of soil water used for growth in early summer. Vigorous stands occur where the precipitation exceeds 533 millimeters (21 inches). These lodgepole forests are typically associated with rock types weathering to acidic substrates, such as granite and rhyolite. In west-central Montana ranges such the Big Belts and the Rocky Mountain Front, these forests are found on limestone substrates. These systems are especially well developed on the broad ridges and high valleys near and east of the Continental Divide. Succession proceeds at different rates, moving relatively quickly on low-elevation, mesic sites and particularly slowly in high-elevation forests such as those along the Continental Divide in Montana.



6% (93,888
Acres)

Grassland Systems Montane Grassland

Rocky Mountain Subalpine-Montane Mesic Meadow

This system is restricted to sites from lower montane to subalpine elevations where finely textured soils, snow deposition, or windswept conditions limit tree establishment. Many occurrences are small patches, and are often found in mosaics within woodlands, dense shrublands, or just below alpine communities. Elevations range from 600 to 2,011 meters (2,000-6,600 feet) in the northern Rocky Mountains and up to 2,286- 2,682 meters (7,500-8,800 feet) in the mountains of southwestern Montana. This system occurs on gentle to moderate-gradient slopes and in relatively moist habitats. Soils are typically seasonally moist to saturated in the spring, but dry out later in the growing season. At montane elevations, soils are usually clays or silt loams, and some occurrences may have inclusions of hydric soils in low, depressional areas. At subalpine elevations, soils are derived a variety of parent materials, and are usually rocky or gravelly with good aeration and drainage, but with a well developed organic layer. Some occurrences are more heavily dominated by grasses, while others are more dominated by forbs. Common grasses include tufted hairgrass (*Deschampsia caespitosa*), showy oniongrass (*Melica spectabilis*), mountain brome (*Bromus carinatus*), blue wildrye (*Elymus glaucus*), awned sedge (*Carex atherodes*), and small wing sedge (*Carex microptera*). Forb dominated meadows usually comprise a wide species diversity which differs from montane to subalpine elevations. Shrubs such as shrubby cinquefoil (*Dasiphora fruticosa* ssp. *floribunda*) and snowberry (*Symphoricarpos* species) are occasional but not abundant. This system differs from the Rocky Mountain Alpine Montane Wet Meadow system in that it soils dry out by mid-summer.



5% (69,806
Acres)

Human Land Use Agriculture

Cultivated Crops

These areas used for the production of crops, such as corn, soybeans, small grains, sunflowers, vegetables, and cotton, typically on an annual cycle. Agricultural plant cover is variable depending on season and type of farming. Other areas include more stable land cover of orchards and vineyards.



2% (26,072
Acres)

Forest and Woodland Systems Conifer-dominated forest and woodland (xeric-mesic)

Rocky Mountain Subalpine Dry-Mesic Spruce-Fir Forest and Woodland

Engelmann spruce (*Picea engelmannii*) and subalpine fir (*Abies lasiocarpa*) make up a substantial part of the montane and lower subalpine forests of the Montana Rocky Mountains and mountain island ranges of north-central and west-central Montana. Spruce is usually associated with fir and occurs as either a climax co-dominant or as a persistent, long-lived seral species in most upper elevation fir habitat types. Dry to mesic spruce-dominated forests range from 884-1,585 meters (2,900-5,200 feet) west of the Continental Divide, and 1585-2,073 meters (5,200-6,800 feet) east of the Continental Divide in the northern and central portions of the state. This system can be found at elevations up to 2,896 meters (9,500 feet) in southwestern Montana. Forests are found on gentle to very steep mountain slopes, high-elevation ridge tops and upper slopes, plateau-like surfaces, basins, alluvial terraces, well-drained benches, and inactive stream terraces. Tree canopy characteristics are relatively uniform. In northern Montana, Engelmann spruce hybridizes with its boreal counterpart, white spruce (*Picea glauca*). Douglas-fir (*Pseudotsuga menziesii*), lodgepole pine (*Pinus contorta*), and western larch (*Larix occidentalis*) (west of the Continental Divide) are seral but often present in these forests. The understory is comprised of a mixture of shrubs, forbs and graminoids tolerant of warmer and drier soil conditions than those found on the more mesic to wet spruce-fir system. The drier occurrences of this system are especially common on steep slopes at upper elevations throughout the eastern Rocky Mountains, whereas the more mesic occurrences form substantial cover west of the Continental Divide in the Flathead, Lolo, Bitterroot and Kootenai river drainages.



Forest and Woodland Systems
Conifer-dominated forest and woodland (xeric-mesic)

Rocky Mountain Ponderosa Pine Woodland and Savanna

2% (24,879 Acres)

This system occurs on warm, dry, exposed sites in the foothills of the Rocky Mountains in west-central and central Montana, at the ecotone between grasslands or shrublands and more mesic coniferous forests. Elevations range from 1,066 to 1,676 meters (3,500-5,500 feet), with higher elevation examples mostly confined to central Montana. Occurrences are found on all slopes and aspects; however, moderately steep to very steep slopes or ridgetops are most common. True savanna types are infrequent; the system is more characteristically an open forest with a grassy understory. In the western part of the state, this system is seen mostly on dry slopes in the rainshadow of the Bitterroot Mountains. East of the Continental Divide, it is most widespread around Helena and Lewistown, although it occurs throughout mountain ranges as far east as the Little Rocky and Bearpaw Mountains. Ponderosa pine (*Pinus ponderosa*) is the dominant conifer. Douglas-fir (*Pseudotsuga menziesii*) and western larch (*Larix occidentalis*) may be present in the tree canopy in the more western areas, but are usually absent. In central Montana, limber pine (*Pinus flexilis*) and horizontal juniper (*Juniperus horizontalis*) are frequently components. Although the understory of ponderosa pine forests is often shrubby in other states, in Montana, habitats are mostly dominated by graminoids, although bitterbrush (*Purshia tridentata*), white snowberry (*Symphoricarpos albus*), and skunkbrush (*Rhus trilobata*) occur in forests on benchlands and rocky slopes in the central portion of the state. Understory vegetation is more typically grasses and forbs that resprout following low to moderate intensity surface fires. Prolonged drought, beetle kill and exotic invasion are rapidly changing the dynamics of this system.




Wetland and Riparian Systems
Floodplain and Riparian

Rocky Mountain Lower Montane-Foothill Riparian Woodland and Shrubland

2% (23,442 Acres)

This ecological system is found throughout the Rocky Mountain and Colorado Plateau regions. In Montana, it ranges from approximately 945 to 2,042 meters (3,100 to 6,700 feet), characteristically occurring as a mosaic of multiple communities that are tree-dominated with a diverse shrub component. It is dependent on a natural hydrologic regime, especially annual to episodic flooding. Occurrences are found within the flood zone of rivers, on islands, sand or cobble bars, and on immediate streambanks. It can form large, wide occurrences on mid-channel islands in larger rivers or narrow bands on small, rocky canyon tributaries and well-drained benches. It is also typically found in backwater channels and other perennially wet but less scoured sites, such as floodplains swales and irrigation ditches. In some locations, occurrences extend into moderately high intermountain basins where the adjacent vegetation is sage steppe. Dominant trees may include boxelder maple (*Acer negundo*), narrowleaf cottonwood (*Populus angustifolia*), Plains cottonwood (*Populus deltoides*), Douglas-fir (*Pseudotsuga menziesii*), peachleaf willow (*Salix amygdaloides*), or Rocky Mountain juniper (*Juniperus scopulorum*). Dominant shrubs include Rocky Mountain maple (*Acer glabrum*), thinleaf alder (*Alnus incana*), river birch (*Betula occidentalis*), redbud (*Cornus sericea*), hawthorne (*Crataegus spp.*), chokecherry (*Prunus virginiana*), skunkbush sumac (*Rhus trilobata*), Drummond's willow (*Salix drummondiana*), sandbar willow (*Salix exigua*), Pacific willow (*Salix lucida*), rose (*Rosa species*), silver buffaloberry (*Shepherdia argentea*), or snowberry (*Symphoricarpos species*). Exotic trees of Russian olive (*Elaeagnus angustifolia*) and saltcedar (*Tamarix species*) may invade some stands in southeastern and south-central Montana.




Recently Disturbed or Modified
Insect-Killed Forest


Insect-Killed Forest


2% (23,213 Acres)


Additional Limited Land Cover


- 1% (20,584 Acres) [Northern Rocky Mountain Lower Montane Riparian Woodland and Shrubland](#)
- 1% (15,791 Acres) [Aspen Forest and Woodland](#)
- 1% (12,228 Acres) [Rocky Mountain Montane-Foothill Deciduous Shrubland](#)
- 1% (11,898 Acres) [Harvested forest-tree regeneration](#)
- 1% (11,882 Acres) [Rocky Mountain Subalpine-Upper Montane Grassland](#)
- 1% (10,753 Acres) [Other Roads](#)
- 1% (10,053 Acres) [Pasture/Hay](#)
- 1% (9,472 Acres) [Harvested forest-grass regeneration](#)
- <1% (6,867 Acres) [Harvested forest-shrub regeneration](#)
- <1% (5,919 Acres) [Burned Sagebrush](#)
- <1% (5,833 Acres) [Rocky Mountain Foothill Limber Pine - Juniper Woodland](#)
- <1% (4,964 Acres) [Recently burned forest](#)
- <1% (4,878 Acres) [Alpine-Montane Wet Meadow](#)
- <1% (4,699 Acres) [Introduced Upland Vegetation - Annual and Biennial Forbland](#)
- <1% (4,135 Acres) [Rocky Mountain Subalpine Mesic Spruce-Fir Forest and Woodland](#)
- <1% (2,369 Acres) [Major Roads](#)
- <1% (2,160 Acres) [Alpine Bedrock and Scree](#)
- <1% (2,137 Acres) [Rocky Mountain Subalpine Deciduous Shrubland](#)
- <1% (1,872 Acres) [Great Plains Shrubland](#)


<1% (1,764 Acres)  [Open Water](#)


<1% (1,303 Acres)  [Low Intensity Residential](#)


<1% (1,285 Acres)  [Rocky Mountain Subalpine Woodland and Parkland](#)


<1% (816 Acres)  [Rocky Mountain Cliff, Canyon and Massive Bedrock](#)


<1% (756 Acres)  [Great Plains Ponderosa Pine Woodland and Savanna](#)


<1% (703 Acres)  [Great Plains Wooded Draw and Ravine](#)


<1% (494 Acres)  [Post-Fire Recovery](#)


<1% (454 Acres)  [Developed, Open Space](#)


<1% (438 Acres)  [Recently burned shrubland](#)


<1% (349 Acres)  [Great Plains Mixedgrass Prairie](#)


<1% (340 Acres)  [Low Sagebrush Shrubland](#)


<1% (252 Acres)  [Alpine Fell-Field](#)


<1% (249 Acres)  [Aspen and Mixed Conifer Forest](#)


<1% (195 Acres)  [Rocky Mountain Foothill Woodland-Steppe Transition](#)


<1% (172 Acres)  [Emergent Marsh](#)


<1% (84 Acres)  [Mountain Mahogany Woodland and Shrubland](#)


<1% (72 Acres)  [Commercial / Industrial](#)


<1% (57 Acres)  [Rocky Mountain Subalpine-Montane Riparian Woodland](#)


<1% (53 Acres)  [Great Plains Badlands](#)


<1% (49 Acres)  [Recently burned grassland](#)


<1% (48 Acres)  [Great Plains Cliff and Outcrop](#)


<1% (44 Acres)  [High Intensity Residential](#)


<1% (29 Acres)  [Rocky Mountain Subalpine-Montane Riparian Shrubland](#)


<1% (6 Acres)  [Quarries, Strip Mines and Gravel Pits](#)


<1% (6 Acres)  [Greasewood Flat](#)


<1% (5 Acres)  [Rocky Mountain Poor Site Lodgepole Pine Forest](#)


<1% (4 Acres)  [Rocky Mountain Conifer Swamp](#)


<1% (2 Acres)  [Great Plains Closed Depressional Wetland](#)

<1% (1 Acres)  [Wind Turbine](#)

<1% (1 Acres)  [Great Plains Sand Prairie](#)

<1% (0 Acres)  [Great Plains Saline Depression Wetland](#)

<1% (0 Acres)  [Rocky Mountain Subalpine-Montane Fen](#)

<1% (0 Acres)  [Rocky Mountain Wooded Vernal Pool](#)

Introduction to Land Cover

Land Use/Land Cover is one of 15 [Montana Spatial Data Infrastructure](#) framework layers considered vital for making statewide maps of Montana and understanding its geography. The layer records all Montana natural vegetation, land cover and land use, classified from satellite and aerial imagery, mapped at a scale of 1:100000, and interpreted with supporting ground-level data. The baseline map is adapted from the Northwest ReGAP (NWGAP) project land cover classification, which used 30m resolution multi-spectral Landsat imagery acquired between 1999 and 2001. Vegetation classes were drawn from the Ecological System Classification developed by NatureServe (Comer et al. 2003). The land cover classes were developed by Anderson et al. (1976). The NWGAP effort encompasses 12 map zones. Montana overlaps seven of these zones. The two NWGAP teams responsible for the initial land cover mapping effort in Montana were Sanborn and NWGAP at the University of Idaho. Both Sanborn and NWGAP employed a similar modeling approach in which Classification and Regression Tree (CART) models were applied to Landsat ETM+ scenes. The Spatial Analysis Lab within the Montana Natural Heritage Program was responsible for developing a seamless Montana land cover map with a consistent statewide legend from these two separate products. Additionally, the Montana land cover layer incorporates several other land cover and land use products (e.g., MSDI Structures and Transportation themes and the Montana Department of Revenue Final Land Unit classification) and reclassifications based on plot-level data and the latest NAIP imagery to improve accuracy and enhance the usability of the theme. Updates are done as partner support and funding allow, or when other MSDI datasets can be incorporated. Recent updates include fire perimeters and agricultural land use (annually), energy developments such as wind, oil and gas installations (2014), roads, structures and other impervious surfaces (various years): and local updates/improvements to specific ecological systems (e.g., central Montana grassland and sagebrush ecosystems). Current and previous versions of the Land Use/Land Cover layer with full metadata are available for download at the Montana State Library's [Geographic Information Clearinghouse](#).

Within the report area you have requested, land cover is summarized by acres of Level 1, Level 2, and Level 3 Ecological Systems.

Literature Cited

- Anderson, J.R. E.E. Hardy, J.T. Roach, and R.E. Witmer. 1976. A land use and land cover classification system for use with remote sensor data. U.S. Geological Survey Professional Paper 964.
- Comer, P., D. Faber-Langendoen, R. Evans, S. Gawler, C. Josse, G. Kittel, S. Menard, M. Pyne, M. Reid, K. Schulz, K. Snow, and J. Teague. 2003. Ecological systems of the United States: A working classification of U.S. terrestrial systems. NatureServe, Arlington, VA.