

County	Yellowstone	Upstream River Mile	331.8
Classification	PCA: Partially confined anabranching	Downstream River Mile	322.7
General Location	Bull Mountain	Length	9.10 mi (14.65 km)

Narrative Summary

Reach B8 is located downstream of Pompey's Pillar. The Reach is 9.1 miles long and is partially confined by the valley wall with numerous forested islands. In the 1950's, the main channel flowed more closely along the north valley wall; southward migration since that time has reduced the influence of the valley wall on stream geomorphology. The valley is wide in this area, which is typical where the bounding rock units are made up of the relatively erodible Cretaceous-age Bearpaw shale.

Just over 3,000 feet of streambank are armored by rock riprap, which is about 3.3 percent of the total bankline. All of the bank armor in the reach is protecting the rail line on the south side of the river. High resolution imagery from fall 2011 indicates that at RM 328 about 570 feet of rock riprap has been flanked on the right bank against the rail line, and that the flanked rock is about 80 feet into the river off of the south bank. Currently, the river is within 100 feet of the rail line and migrating rapidly in that direction.

One side channel that is about 6,200 feet long at RM 326R was blocked prior to 1950.

Land uses in the reach are primarily agricultural, with about 1,240 acres of flood irrigated land mapped as of 2011. There are 124 acres of land in sprinkler and 86 under pivot. The modern 5-year floodplain contains about 250 acres of flood-irrigated ground.

One dump site was mapped on an old swale adjacent to a flood irrigated field at RM 326.5R.

The Channel Migration Zone (CMZ) has been developed for primarily flood irrigation; as of 2011, there were 457 acres of flood irrigated land in the CMZ, and about 7 percent of the total CMZ footprint has become restricted by bank armor and road prisms. The railroad has isolated almost 9 percent of the historic 100-year floodplain in the reach. About 22 percent of the 5-year floodplain has become isolated in Reach B8. Much of that 5-year floodplain isolation is due to transportation infrastructure on the south side of the river.

Similar to Reach B7 upstream, Reach B8 shows major southward migration of the river since 1950, with one area at RM 324.3 experiencing over 1,500 feet of migration over the past 60 years. This southward migration has threatened the rail line at RM 328R.

Overall, the migration rates and floodplain turnover rates have dropped since 1976 from 1.9 acres/valley mile/year from 1950 to 1976 to 1.5 acres/valley mile/year from 1976-2001.

Reach B8 has 91 mapped acres of Russian olive that can be found in dense stands, especially on forested islands. Even so, the extensive lateral migration of the river has promoted extensive recruitment of new woody riparian habitat. Since the 1950s there has been about 600 acres of riparian recruitment in the reach, most of which was riparian colonization of old 1950's channel area. The acreage of recruitment has exceeded that of erosion of riparian areas by 51 acres. Additionally, there are 271 mapped wetlands in the reach, including 147 acres of wet meadows and marsh. The reach contains about 33 wetland acres per valley mile, which is a relatively high value for the Yellowstone River.

Reach B8 was sampled as part of the avian study. The average species richness in this reach was 7.8, which indicates the average number of species observed during site visits to the reach in cottonwood habitats. The average species richness for sites evaluated is 8. One bird species identified by the Montana Natural Heritage Program as a Potential Species of Concern was identified, the Plumbeous Vireo. Another species identified as a Species of Concern was identified, the Red-headed Woodpecker.

A hydrologic evaluation of flow depletions indicates that flow alterations over the last century have been major in this reach. The mean annual flood is estimated to have dropped from 28,000 cfs to 22,800 cfs, a drop of about 19 percent. The 2-year flood, which strongly influences overall channel form, has dropped by 11 percent. Low flows have also been impacted; severe low flows described as 7Q10 (the lowest average 7-day flow anticipated every ten years) for summer months has dropped from an estimated 3,040 cfs to 2,070 cfs with human development, a reduction of 32 percent. More typical summer low flows, described as the summer 95% flow duration, have dropped from 3,846 cfs under unregulated conditions to 2,227 cfs under regulated conditions at the Billings gage, a reduction of 42 percent.

CEA-Related observations in Reach B8 include:

- Migration away from valley wall resulting in loss of bluff pool habitat.
- Blockage of one side channel at RM 326 sometime prior to 1950
- Transportation infrastructure –caused isolation of 5-year floodplain south of the river at RM 329.5


Recommended Practices (may include Yellowstone River Recommended Practices--YRRPs) for Reach B8 include:

- Side channel reactivation at RM 326
- Dump removal at RM 326.5R
- Flanked armor removal at RM 328R
- Russian olive removal

The following table summarizes some key CEA results that have been used to describe overall condition and types of human influences affecting the river. The values are specific to this single reach. Blanks indicate that a particular value was not available for this area. This information is consolidated from a large dataset that is presented in more detail in the full reach narrative report.

Discharge	Undev.	Developed	% Change	"Undeveloped" flows represent conditions prior to significant human development, whereas "developed" flows reflect the current condition of both consumptive and non-consumptive water use.		
2 Year (cfs)	51,700	46,100	-10.8%			
100 Year (cfs)	90,900	87,600	-3.6%			
Bankfull Channel Area (Ac)	1950	1976	1995	2001	1950-2001	Bankfull channel area is the total footprint of the river inundated at approx. the 2-year flood.
	1,051.1	1,093.5	1,003.0	1,089.4	38.3	
Physical Features	2011 Length (ft)	% of Bankline	2001-2011 Change	There are additional types of bank armor such as car bodies and steel retaining walls, but they are relatively minor.		
Rock RipRap	3,208	3.3%	0			
Concrete Riprap	0	0.0%	0			
Flow Deflectors	0	0.0%	0			
Total	3,208	3.3%	0			
Length of Side Channels Blocked (ft)	Pre-1950s	Post-1950s	Numerous side channels have been blocked by small dikes.			
	6,209	0				
Floodplain Turnover	1950 - 1976	1976 - 2001	1950-2001 In-channel riparian encroachment (negative number indicates retreat)		The rate of floodplain turnover reflects how many acres of land are eroded by the river. Turnover is associated with the creation of riparian habitat.	
Total Acres	391.0	291.8	50.51 acres			
Acres/Year	15.0	11.7				
Acres/Year/Valley Mile	1.9	1.5				
Open Bar Area	Point Bars	Bank Attached	Mid-Channel	Total	The type and extent of open sand and gravel bars reflect in-stream habitat conditions that can be important to fish, amphibians, and ground-nesting birds such as least terns.	
Change in Area '50 - '01 (Ac)						
Floodplain Isolation	Acres	% of FP	Floodplain isolation refers to area that historically was flooded, but has become isolated do to flow alterations or physical features such as levees.			
5 Year	442.3	22%				
100 Year	219.4	9%				
Restricted Migration Area	Acres	% of CMZ	Channel Migration Zone restrictions refer to the area and percent of the CMZ that has been isolated by features such as bank armor, dikes, levees, and transportation embankments.			
	224.3	7%				
Land Use	1950	2011	1950	2011	Changes in land use reflect the development of the river corridor through time. The irrigated agricultural are is a sub-set of the mapped agricultural land.	
Agricultural Land (Ac)	4,889.1	4,506.4	Flood (Ac)	1,269.7	1,238.8	
Ag. Infrastructure (Ac)	90.7	123.1	Sprinkler (Ac)	6.1	124.4	
Exurban (Ac)	43.0	77.4	Pivot (Ac)	0.0	85.9	
Urban (Ac)	0.0	0.0				
Transportation (Ac)	105.3	235.1				
1950s Riparian Vegetation Converted to a Developed Land Use (ac)	To Irrigated	To Other Use	Total Rip. Converted	% of 1950s Rip.	Changes in the extents of riparian vegetation are influenced by land use changes within the corridor.	
	46.9	0.0	46.9	4.0%		
National Wetlands Inventory	Acres	Acres per Valley Mi	Total Wetland Acres	Wetlands units summarized from National Wetlands Inventory Mapping include Riverine (typically open water sloughs), Emergent (marshes and wet meadows) and Shrub-Scrub (open bar areas with colonizing woody vegetation).		
Riverine	10.3	1.3	271.4			
Emergent	147.4	18.8				
Scrub/Shrub	113.7	14.5				
Russian Olive (2001) (Appx. 100-yr Floodplain)	Acres	%	Russian olive is considered an invasive species and its presence in the corridor is fairly recent. Its spread can be used as a general indicator of invasive plants within the corridor.			
	91.2	3.2%				
Riparian Forest at low risk of Cowbird Parasitism (Ac/Valley Mile)	1950	1976	2001	Change 1950-2011	Cowbirds are associated with agricultural and residential development, displacing native bird species by parasitizing their nests.	
	8.5	7.5	8.4	0.0		

PHYSICAL FEATURES MAP (2011)



CHANNEL MIGRATION ZONE MAP

