

County	Richland	Upstream River Mile	49.9
Classification	PCA: Partially confined anabranching	Downstream River Mile	36.3
General Location	Seven Sisters	Length	13.60 mi (21.89 km)

Narrative Summary

Reach D12 is located in Richland County at Seven Sisters. The Seven Sisters Fishing Access Site is located in the lower portion of the reach. The reach is a 13.6 mile long Partially Confined Anabranching reach type, indicating some influence of the valley wall along with extensive forested islands. This reach supports over 20 miles of side channels, and islands that are miles long and over ½ mile wide.

There are almost 7,000 feet of bank armor in the reach, and about one third of that was built since 2001. Most of the armor (3,250 feet) is rock riprap, and there are about 2,000 feet each of concrete riprap and flow deflectors. A total of 5 percent of the bank is armored, which is a relatively low concentration of bank armor for the Yellowstone River. All of the armor is protecting agricultural land, most of it against a flood irrigated field on the left bank in the lower end of the reach at RM 37.

Since 1950, a side channel that is almost three miles long was blocked at RM 45.3L. There have also been some gains in side channel length in the reach, such that the net change in length is a loss of approximately one mile. As of 2001, this reach supported almost 21 miles of anabranching channel.

Land use is dominated by agriculture, with 583 acres of pivot irrigation development since 1950. Physical features such as bank armor, dikes, and levees have isolated 3 percent of the Channel Migration Zone in Reach D12, and as of 2011 there were 224 acres of land in the CMZ under pivot irrigation, and 900 acres under flood.

Reach D12 shows, like most other reaches below the Bighorn River, a shrinking channel with reduced rates of erosion and floodplain turnover. For example, the bankfull channel area in the reach dropped by 480 acres since 1950, and there was almost 600 acres of riparian encroachment into old channel areas. Floodplain turnover rates have dropped from 2.1 acres/valley mile/year from 1950-1976 to 1.3 acres/valley mile/year from 1976-2001. This equates to 330 fewer acres of floodplain turnover since 1976. There has also been a net loss of 159 acres of open bar area as the channel has become smaller and more forested. On the floodplain, riparian acreage has decreased; about 350 acres or 9 percent of the total riparian area was cleared for irrigation since 1950.

There are 75 acres of Russian olive in the reach.

The 100-year floodplain has been isolated in this reach, but compared to other reaches the isolation has been fairly minor. About 300 acres of 100-year floodplain has been isolated by human development, which is 5 percent of the total 100-year floodplain. Although only about 5 percent of the 100-year floodplain has been isolated, the impact of flow alterations on the smaller 5-year floodplain has been much more severe; 42 percent of the historic 5-year floodplain is no longer inundated at that frequency. The isolation of the historic 5-year floodplain, which is due primarily to flow alterations, has been associated with increased development in these areas; currently there are about 300 acres of flood irrigated land and within the historic 5-year floodplain footprint.

There is an animal feeding facility on the right bank at RM 46.8.

Reach D12 was sampled as part of the fisheries study. A total of 37 fish species were sampled in the reach. Three species collected in the reach have been identified by the Montana Natural Heritage Program as Species of Concern (SOC): Pallid Sturgeon, Sauger, and Sturgeon Chub.

Reach D12 was also sampled as part of the avian study. A total of 59 bird species were identified in the reach. All five bird species identified by the Montana Natural Heritage Program as Potential Species of Concern (PSOC) on the Yellowstone River were also found, the Black and White Warbler, the Chimney Swift, the Dickcissel, the Ovenbird, and the Plumbeous Vireo. Similarly, all three bird species identified as Species of Concern (SOC) were identified: the Black-billed Cuckoo, Bobolink, and Red-headed Woodpecker. In contrast to most other reaches, Reach D12 has seen an increase in the forested area that is at low risk of cowbird parasitism since 1950. At that time, there were 103 acres per valley mile of such forest, and that number increased to 115 acres per valley mile by 2001.

A hydrologic evaluation of flow depletions indicates that flow alterations over the last century have been major in this reach. The 2-year flood, which strongly influences overall channel form, has dropped by 22 percent. Low flows have also been impacted; severe low flows described as 7Q10 (the lowest average 7-day flow anticipated every ten years) for summer months has dropped from an estimated 4,310 cfs to 2,410 cfs with human development, a reduction of 50 percent. More typical summer low flows, described as the summer 95% flow duration, have dropped from 6,470 cfs under unregulated conditions to 2,680 cfs under regulated conditions, a reduction of 59 percent.

CEA-Related observations in Reach D12 include:

- Increase in area at low risk of cowbird parasitism with riparian encroachment


Recommended Practices (may include Yellowstone River Recommended Practices--YRRPs) for Reach D12 include:

- Nutrient management at animal handling facility at RM 46.8R
- Side channel reactivation at RM 45.3R
- Russian olive removal

The following table summarizes some key CEA results that have been used to describe overall condition and types of human influences affecting the river. The values are specific to this single reach. Blanks indicate that a particular value was not available for this area. This information is consolidated from a large dataset that is presented in more detail in the full reach narrative report.

Discharge	Undev.	Developed	% Change	"Undeveloped" flows represent conditions prior to significant human development, whereas "developed" flows reflect the current condition of both consumptive and non-consumptive water use.		
2 Year (cfs)	69,800	54,300	-22.2%			
100 Year (cfs)	144,000	132,000	-8.3%			
Bankfull Channel Area (Ac)	1950	1976	1995	2001	1950-2001	Bankfull channel area is the total footprint of the river inundated at approx. the 2-year flood.
	2,239.4	1,957.5	1,919.3	1,754.7	-484.8	
Physical Features	2011 Length (ft)	% of Bankline	2001-2011 Change	There are additional types of bank armor such as car bodies and steel retaining walls, but they are relatively minor.		
Rock Riprap	3,251	2.3%	2,655			
Concrete Riprap	1,945	1.4%	0			
Flow Deflectors	1,801	1.3%	118			
Total	6,997	4.9%	2,773			
Length of Side Channels Blocked (ft)	Pre-1950s	Post-1950s	Numerous side channels have been blocked by small dikes.			
	0	14,624				
Floodplain Turnover	1950 - 1976	1976 - 2001	1950-2001 In-channel riparian encroachment (negative number indicates retreat)		The rate of floodplain turnover reflects how many acres of land are eroded by the river. Turnover is associated with the creation of riparian habitat.	
Total Acres	596.0	338.4	597.01 acres			
Acres/Year	22.9	13.5				
Acres/Year/Valley Mile	2.1	1.3				
Open Bar Area	Point Bars	Bank Attached	Mid-Channel	Total	The type and extent of open sand and gravel bars reflect in-stream habitat conditions that can be important to fish, amphibians, and ground-nesting birds such as least terns.	
Change in Area '50 - '01 (Ac)	-205.7	27.4	19.8	-158.5		
Floodplain Isolation	Acres	% of FP	Floodplain isolation refers to area that historically was flooded, but has become isolated do to flow alterations or physical features such as levees.			
5 Year	2,113.3	42%				
100 Year	344.5	5%				
Restricted Migration Area	Acres	% of CMZ	Channel Migration Zone restrictions refer to the area and percent of the CMZ that has been isolated by features such as bank armor, dikes, levees, and transportation embankments.			
	197.9	3%				
Land Use	1950	2011	1950	2011	Changes in land use reflect the development of the river corridor through time. The irrigated agricultural are is a sub-set of the mapped agricultural land.	
Agricultural Land (Ac)	5,885.9	6,086.8	Flood (Ac)	2,107.6	2,364.7	
Ag. Infrastructure (Ac)	59.8	154.9	Sprinkler (Ac)	0.0	0.0	
Exurban (Ac)	0.0	1.7	Pivot (Ac)	0.0	582.7	
Urban (Ac)	0.0	0.0				
Transportation (Ac)	43.7	58.6				
1950s Riparian Vegetation Converted to a Developed Land Use (ac)	To Irrigated	To Other Use	Total Rip. Converted	% of 1950s Rip.	Changes in the extents of riparian vegetation are influenced by land use changes within the corridor.	
	353.9	0.8	354.7	9.0%		
National Wetlands Inventory	Acres	Acres per Valley Mi	Total Wetland Acres	Wetlands units summarized from National Wetlands Inventory Mapping include Riverine (typically open water sloughs), Emergent (marshes and wet meadows) and Shrub-Scrub (open bar areas with colonizing woody vegetation).		
Riverine	28.0	2.6	285.0			
Emergent	117.2	10.9				
Scrub/Shrub	139.8	13.0				
Russian Olive (2001) (Appx. 100-yr Floodplain)	Acres	%	Russian olive is considered an invasive species and its presence in the corridor is fairly recent. Its spread can be used as a general indicator of invasive plants within the corridor.			
	74.8	1.4%				
Riparian Forest at low risk of Cowbird Parasitism (Ac/Valley Mile)	1950	1976	2001	Change 1950-2011	Cowbirds are associated with agricultural and residential development, displacing native bird species by parasitizing their nests.	
	103.2	104.8	115.5	12.3		

PHYSICAL FEATURES MAP (2011)



CHANNEL MIGRATION ZONE MAP

